



Pericardial Manifestations in Autoimmune Encounters: Some Worthy Facts for Consideration!

Lovely Chhabra, Nauman Khalid¹, David H. Spodick²

Department of Cardiovascular Medicine, Hartford Hospital, University of Connecticut, School of Medicine, Hartford, CT, ¹Department of Internal Medicine, Hartford Hospital, University of Connecticut, School of Medicine, Hartford, CT, ²Department of Cardiovascular Medicine, Saint Vincent Hospital, University of Massachusetts Medical School, Worcester, MA, USA

Date of Submission: May 20, 2014

Date of Acceptance: Sep 27, 2014

Correspondence to:

Dr. Lovely Chhabra,
80, Seymour Street, Hartford,
CT 06102, USA.
E-mail: lovids@hotmail.com

How to cite this article: Chhabra L, Khalid N, Spodick DH. Pericardial Manifestations in Autoimmune Encounters: Some Worthy Facts for Consideration!. Int J Prev Med 2014;5:1632-4.

DEAR EDITOR,

We read with great pleasure the manuscript by Ashrafi *et al.* published in a recent issue of the International Journal of Preventive Medicine.^[1] Their case highlights a very interesting association between celiac disease and pericardial manifestations. However, authors have not discussed the underlying mechanism for its causal association and treatment considerations for refractory cases.

Pericarditis has previously been reported in celiac disease, and it may even manifest as a primary presenting feature.^[2] The likely mechanism is a disturbed humoral and cell-mediated immunity including other autoimmune reactions. There is deposition of circulating immune complexes originating from the small intestine, which may resemble the etiology of pericarditis similar to the one encountered in serum sickness.^[2] The mechanisms are likely related to pericardial deposition of soluble antigen-antibody complexes when there is excess of antigen. It may often be difficult to differentiate the pericardial syndromes encountered in inflammatory bowel disease than those with celiac disease as the clinical picture is often similar. However, the response to gluten free diet often makes the etiological association very

convincing.^[2] Gluten free diet thus not only serves a diagnostic clinical marker, but also an effective treatment modality for this syndrome.^[2]

We also want to highlight that celiac disease may be associated with several autoimmune conditions such as hypothyroidism, adrenal sufficiency, or polyglandular endocrine insufficiency syndromes etc., which can in fact independently contribute to the underlying etiopathogenesis of generalized serositis including pericardial effusion.^[2] In our clinical experience, we have had encountered patients with polyglandular insufficiency who presented with pericardial effusion and even tamponade, in the setting of thyroid and adrenal deficiency. Thus, we recommend that clinicians should be vigilant about their coexistence and an appropriate screening for these conditions should be sought under appropriate clinical settings for example if patients have not responded to the standard treatment therapy of celiac disease. Authors have commendably excluded hypothyroidism in the presented case. This differential consideration is important for the above-mentioned reasons. After appropriately excluding other autoimmune conditions, if pericardial symptoms of celiac disease patients do not completely respond to gluten free diet alone, then steroids should be considered as a

first-line therapy.^[3] It is in contrast to the treatment of acute idiopathic pericarditis where one would consider either nonsteroidal antiinflammatory drugs or colchicine.^[3] Colchicine has not been extensively studied in patients with autoimmune diseases; however at least 7% of the patients in CORP-2 trial had underlying autoimmune conditions and benefited from the use of colchicine.^[4] Thus, addition of colchicine to steroids therapy may be considered in such patients though extensive prospective data to support this recommendation is lacking.

Clinicians should be aware of these indications in order to make an appropriate informed decision and selection of treatment therapy.

REFERENCES

1. Ashrafi F, Darakhshandeh A, Heidarpour M, Tavakoli T,

- Najafian J. Pericardial effusion in celiac disease. *Int J Prev Med* 2014;5:356-9.
2. Spodick DH. *The Pericardium: A Comprehensive Textbook*. New York: Marcel Dekker; 1997. p. 330-1.
 3. Chhabra L, Spodick DH. Letter by Chhabra and Spodick regarding article, "Treatment of acute and recurrent idiopathic pericarditis". *Circulation* 2013;128:e391.
 4. Imazio M, Belli R, Brucato A, Cemin R, Ferrua S, Beqaraj F, *et al.* Efficacy and safety of colchicine for treatment of multiple recurrences of pericarditis (CORP-2): A multicentre, double-blind, placebo-controlled, randomised trial. *Lancet* 2014;383:2232-7.

Source of Support: Nil, **Conflict of Interest:** Dr. David H. Spodick receives royalties from his textbook, "The Pericardium: A Comprehensive Textbook (Fundamental and Clinical Cardiology), Marcel Dekker, New York 1997."

International Journal of Preventive Medicine on Web

<http://www.journalonweb.com/ijpm>

International Journal of Preventive Medicine now accepts articles electronically. It is easy, convenient and fast. Check following steps:

1 Registration

- Register from <http://www.journalonweb.com/ijpm> as a new author (Signup as author)
- Two-step self-explanatory process

2 New article submission

- Prepare your files (Article file, First page file and Images, if any)
- Login into your area
- Click on 'Submit a new article' under 'New Article'
- Follow the steps (three steps for article without images and five for with images)
- On successful submission you will receive an acknowledgement quoting the manuscript numbers

3 Tracking the progress

- Click on 'In Review Article' under 'Submitted Articles'
- The table gives status of the article and its due date to move to next phase
- More details can be obtained by clicking on the Manuscript ID
- Comments sent by the editor and referee will be available from these pages

4 Submitting a revised article

- Click on 'Article for Revision' under 'Submitted Articles'
- Click on 'Revise'
- From the first window, you can modify Article Title, Article Type
- First Page file and Images could be modified from second and third window, respectively
- The fourth step is uploading the revised article file.
- Include the referees' comments along with the point to point clarifications at the beginning of the revised article file.
- Do not include authors' name in the article file.
- Upload the revised article file against New Article File - Browse, choose your file and then click "Upload" OR Click "Finish"
- On completion of revision process you will be able to check the latest file uploaded from Article Cycle (In Review Articles-> Click on manuscript ID -> Latest file will have a number with 'R')

Facilities

- Submission of new articles with images
- Submission of revised articles
- Checking of proofs
- Track the progress of article in review process

Advantages

- Any-time, any-where access
- Faster review
- Cost saving on postage
- No need for hard-copy submission (except on acceptance images should be sent)
- Ability to track the progress
- Ease of contacting the journal

Requirements for usage

- Computer and internet connection
- Web-browser (preferably newer versions - IE 5.0 or NS 4.7 and above)
- Cookies and javascript to be enabled in web-browser

Online submission checklist

- First Page File (text/rtf/doc/pdf file) with title page, covering letter, acknowledgement, etc.
- Article File (text/rtf/doc/pdf file) - text of the article, beginning from Title, Abstract till References (including tables). File size limit 1 MB. Do not include images in this file.
- Images (jpeg): Submit good quality colour images. Each image should be less than 4096 kb (4 MB) in size.

Help

- Check Frequently Asked Questions (FAQs) on the site
- In case of any difficulty contact the editor