

The Effect of Hydro-alcoholic Extract of *Rheum Turkestanicum* Roots against Oxidative Stress in Endothelial Cells

Abstract

Introduction: Cardiovascular disorders (CVD) are a common cause of mortality worldwide. Oxidative stress is thought to be a major factor leading to CVD. Anti-oxidants such as medicinal plants may have a role in the mitigation of vascular problems through free radicals scavenging. In this study, we evaluated the protective effects of *Rheum turkestanicum* against hydrogen peroxide (H_2O_2)-induced toxicity in endothelial cells (BAE-1). **Methods:** To evaluate the protective effect of *R. turkestanicum* against H_2O_2 toxicity, four groups comprised of control group (the cells without any treatment), H_2O_2 group (the cells incubated with H_2O_2 (200 μM)), and treatment groups (the cells treated with *R. turkestanicum* (12200 $\mu g/ml$) alone or 24h before exposure to H_2O_2). Quercetin (30.23 $\mu g/ml$) was used as a bioactive ingredient of the extract. Then the cell viability, reactive oxygen species, lipid peroxidation, and apoptosis were evaluated. **Results:** H_2O_2 exposure reduced cell viability to $13.6 \pm 1.6\%$, enhanced ROS generation to $1445 \pm 80.7\%$, lipid peroxidation (LPO, $290 \pm 13\%$ of control), and apoptotic cells ($P < 0.001$). In contrast, compared with H_2O_2 group, *R. turkestanicum* and quercetin significantly restored the cell viability to 80.3 ± 1.6 and $87.2 \pm 2.1\%$, ROS formation to 186 ± 10 and $129 \pm 1\%$, as well as LPO to 130.7 ± 7.7 and 116 ± 2.5 of control, respectively ($P < 0.001$). Therefore, the extract reduced H_2O_2 -induced toxicity in BAE-1 cells by scavenging of free radicals. **Conclusion:** Our findings demonstrated that the extract might reduce toxicity of endothelial cells by attenuation of oxidative stress, which can be related to the presence of active ingredients including quercetin.

Keywords: Apoptosis, endothelial cells, oxidative stress, quercetin, *Rheum turkestanicum*

Introduction

Endothelial cells are involved in vascular hemostasis, angiogenesis, inflammation responses, and vascular contraction.^[1,2] Studies have shown that dysfunction of endothelial cells contributes to the pathogenesis of cardiovascular diseases (CVD).^[2] Most of the risk factors associated with CVD disturb the cell function and processes like apoptosis.^[3] Oxidative stress has been suggested as the common characteristic of risk factors of CVD including hypertension, aging, obesity, unhealthy diet, and low-physical activity.^[3]

Oxidative stress is induced by production of excessive amount of ROS and exhausted antioxidant defense systems. ROS cause DNA injury and mitochondrial dysfunction which consequently results in apoptotic cell death.^[4]

Endothelial dysfunction during atherosclerosis is associated with the release

of arachidonic acid and inflammatory cytokines, including tumor necrosis factor- α (TNF- α), Interleukin-1 (IL-1) and IL-6, and induces apoptosis of endothelial and vascular smooth muscle cells. Inflammation can induce oxidative stress, which in turn can inflammation.^[3,5]

Among different types of ROS, hydrogen peroxide (H_2O_2) is common, which is used extensively in *in vitro* studies to induce endothelial cell injuries.^[6] Although the production of free radicals plays a role in the pathogenesis of different disorders, however, medicinal herbs with anti-oxidant properties such as polyphenols, beta-carotene, and tocopherols may reduce H_2O_2 -induced toxicity in endothelial cells.^[3] Recent studies have reported *Crocus sativus*,^[7] *Phyllanthus emblica*,^[8] *Melissa officinalis*,^[9] and pomegranate seed oil diminished H_2O_2 toxicity through scavenging of free radicals.^[10] *R. turkestanicum* belongs to *Polygonaceae* family and grows in north-east and central Asia particularly

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Hosseini A, Sheikh S, Soukhtanloo M, Malaekheh-Nikouei B, Rajabian A. The effect of hydro-alcoholic extract of *Rheum turkestanicum* roots against oxidative stress in endothelial cells. *Int J Prev Med* 2020;11:122.

Azar Hosseini^{1,2},
Sahar Sheikh²,
Mohammad
Soukhtanloo³,
Bizhan Malaekheh-
Nikouei⁴,
Arezoo Rajabian¹

¹Pharmacological Research Center of Medicinal Plants, Mashhad University of Medical Sciences, Mashhad, Iran,

²Department of Pharmacology, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran, ³Department of Medical Biochemistry, Faculty of Medicine, Mashhad University of Medical Sciences, Mashhad, Iran, ⁴Nanotechnology Research Center, Institute of Pharmaceutical Technology, Mashhad University of Medical Sciences, Mashhad, Iran

Address for correspondence:

Dr. Arezoo Rajabian,
Pharmacological Research Center of Medicinal Plants, Mashhad University of Medical Sciences, Mashhad, Iran.
E-mail: rajabianar@mums.ac.ir

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.net

DOI:
10.4103/ijpvm.IJPVM_386_19

Quick Response Code:



of Iran. Based on the recent pharmacological studies, *R. turkestanicum* has been used as an anticancer, antidiabetic, and antihypertensive.^[11] Recent studies have revealed a protective effect of this plant against toxic agents, such as doxorubicin,^[12] cisplatin,^[13] gentamicin,^[14] mercuric chloride,^[15] and hexachlorobutadien.^[16] Antioxidant activity of some plants of *Rheum* genus has been attributed to the presence of bioactive components that scavenge free radicals (IC₅₀ value in DPPH assay: 2.8-11.8 μM).^[17] *R. turkestanicum* counter acted hexachlorobutadien and mercuric chloride-induced nephrotoxicity through its antioxidant properties indicated by attenuated lipid peroxidation and increased total thiol content. The antioxidant activity of the plant was also involved in the neuroprotective and anti-apoptotic effects that have been linked to the suppression of ROS generation and lipid peroxidation.^[18]

Most of the pharmacological effects may be attributable to the active compounds such as polyphenolic and flavonoid. These bioactive components have found to possess antioxidant and free radicals scavenging potential.^[19] Quercetin as a bioactive ingredient is found in *Rheum* species including *R. turkestanicum*.^[20] In this investigation, we evaluated the protective effects of *R. turkestanicum* and quercetin against H₂O₂-induced toxicity in bovine aortic endothelium (BAE-1) cells.

Materials and Methods

Reagents

3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl tetrazolium (MTT), 2',7'-dichlorofluorescein diacetate (DCFH-DA), propidium iodide (PI), H₂O₂, Triton X-100, thiobarbituric acid (TBA), sodium citrate and quercetin were provided from Sigma. Dulbecco's Modified Eagles Medium (DMEM), fetal bovine serum (FBS), and penicillin/streptomycin ×100 were obtained from Gibco. Dimethyl sulfoxide (DMSO) was purchased from Merck. BAE-1 cell line was prepared from the Pasteur Institute (Tehran, Iran).

Preparation of extract

R. turkestanicum roots were collected from the Kalat region (Khorasan Razavi, Iran). This plant was identified by M.R. Joharchi, Ferdowsi University of Mashhad Herbarium (voucher specimen No. 21377). The roots were dried and crushed into a powder by electric micronizer. The soxhlet extract was prepared by 70% ethanol, then dried and kept at -20°C until downstream processes.

Cell culture

BAE-1 cells were cultured in DMEM supplemented with 10% FBS, 100 μg/ml penicillin and streptomycin at 37°C in a humidified atmosphere (90%) containing 5% CO₂.

Assessment of cell viability

The cell viability was evaluated by MTT assay. The cells were seeded, pre-treated with extract (12200 μg/ml) and quercetin (30.23 μg/ml) for 24 h. After 24 h of incubation, the cells were exposed to H₂O₂ at a dose of 200 μM for 30 min (the concentration was chosen based on the previous study).^[12,21] Then, the MTT solution was added to each well and incubated for 3h, DMSO was used to dissolve the Formazan precipitate. The absorbance of samples was determined at 600 nm on an ELISA reader.^[18]

Measurement of reactive oxygen species

DCFH-DA method was used to determine the intracellular ROS level. The cells were pretreated with the extract (12200 μg/ml) and quercetin (30.23 μg/ml) for 24 h, then H₂O₂ (200 μM) was added to cells for 30 min.^[12] After 30 min exposure to H₂O₂, the cells were treated with DCFH-DA and incubated for 30 min. Fluorescence strength was measured by a microplate reader at 504 nm for excitation and 524 nm for emission.^[22]

Lipid peroxidation assay

ROS can destroy membrane lipids and produce a variety of breakdown products including alcohols, ketones, aldehydes, and others. Malondialdehyde (MDA) is a main product of lipid peroxidation (LPO). MDA reacts with thiobarbituric acid (TBA) to generate fluorescence adduct. LPO was measured by TBA reactive substance (TBARS) assay. The amount of LPO was estimated by TBARS fluorescence intensity.^[23] After the incubation, the cells were scraped and centrifuged at 13,000 ×g for 30 min at 4°C.^[23] Then, 400 μl of trichloroacetic acid (15%) and 800 μl of TBA (0.7%) were added to 500 μl of cells. The mixture was vortexed and then heated for 40 min in a boiling water bath. Subsequently, 200 μl of the sample was transferred to 96-well plate, and the fluorescence intensity was read with excitation and emission of 480 and 530 nm, respectively. The experiment was carried out in triplicate.

Determination of apoptosis

PI-staining was used to identify the apoptotic cells. A sub-G1 peak, indicative of DNA fragmentation, is observed after cell incubation in a hypotonic phosphate-citrate buffer, containing a DNA-binding dye (such as PI). In the histogram, DNA-free apoptotic cells absorb less stain, which can be observed on the left side of the peak. On the basis of the described protocol, the cells were treated after seeding in a 24-well plate. Then, adherent and floating cells were harvested, and incubation was performed at 4°C in darkness overnight, using a hypotonic buffer (750 μl; 50 μg/ml of PI in 0.1% triton X-100 and 0.1% sodium citrate). Finally, a FACS can system (Becton Dickinson) was used to perform flow cytometry, yielding a total of 10⁴ events.^[12]

Statistical analysis

All obtained data were expressed as mean \pm SEM from three independent experiments. Graph Pad Prism version 6 was used to statistically analyze the data. The statistical analysis was performed using Graph Pad Prism version 6. Statistical evaluation was done using one-way analysis of variance, followed by Tukey *post hoc* test. The minimum level of significance was $P < 0.05$.

Results

Effect of hydro-alcoholic extract of *R. turkestanicum* on cell viability

The toxicity effect of *R. turkestanicum* was evaluated on BAE-1 cells at different doses (12200 $\mu\text{g/ml}$) by MTT assay. Results indicated that the extract had no toxic effect on cell viability [Figure 1].

Hydro-alcoholic extract of *R. turkestanicum* attenuated cell death following H_2O_2 toxicity

The results revealed H_2O_2 increased cell death at a dose of 200 μM ($P < 0.001$) compared with the control group, while quercetin and different concentrations of extract (25200 $\mu\text{g/ml}$) reduced cell death and had a protective effect against H_2O_2 -induced toxicity on BAE-1 cells ($P < 0.001$) [Figure 2]. In the cells treated with 100 and 200, the viability significantly increased, compared with that in the groups treated with 12, 25, 50 μM ($P < 0.001$). The cell viability was significantly elevated in the 12 and 100 μM treated groups versus 25 and 200 μM group, respectively ($P < 0.01$, $P < 0.001$). Although, there were significant differences between the extract (12-200 μM) and control group ($P < 0.001$), the viability increased concentration-dependently in the cells treated with the extract. Similarly, significant differences were observed between the 12 and 100 μM treated groups and quercetin ($P < 0.001$).




Figure 1: Effect of hydro-alcoholic extract of *R. turkestanicum* on BAE-1 cell viability. The cells were pretreated with different concentrations of *R. turkestanicum* for 24 h followed by 30 min incubation with H_2O_2 . The cell viability was determined via the MTT assay. The data were expressed as percentage viability of control. The results are mean \pm SEM. The test was done in triplicate ($n = 3$)

Hydro-alcoholic extract of *R. turkestanicum* reduced ROS

Our findings showed that H_2O_2 elevated the level of ROS in the cells ($P < 0.001$). Quercetin and the extract significantly reduced ROS generation in comparison with H_2O_2 group, in a dose-dependently manner [$P < 0.001$, Figure 3]. Compared to the control group, the extract concentrations $< 100 \mu\text{M}$ (50, 25, and 12 μM) exhibited remarkable differences ($P < 0.001$). Intracellular ROS significantly decreased in the cells treated with 100 and 200 μM of the extract, compared with that in the groups treated with 12, 25, and 50 μM of the extract ($P < 0.001$). Therefore, the inhibiting effect of the extract on intracellular ROS was concentration-dependent. ROS content was significantly reduced in the 12 μM treated groups versus 25 μM of the extract ($P < 0.01$), respectively. Similarly, significant differences were observed between the groups treated with the extract at 12100 μM and quercetin ($P < 0.001$).

Hydro-alcoholic extract of *R. turkestanicum* attenuated lipid peroxidation

H_2O_2 increased MDA as lipid peroxidation index ($P < 0.001$). Pretreatment of cells with different doses of extract and quercetin attenuated lipid peroxidation and the level of MDA ($P < 0.001$) [Figure 4]. In the cells treated with the extract at 100 and 200 μM , MDA level significantly decreased, compared with that in the groups treated with 12, 25 μM of the extract ($P < 0.001$). Also, the differences between the cells treated with 50 μM and the cells treated with 12, 25, and 200 μM of the extract were significant ($P < 0.001$). Although, there were significant differences between the extract at different concentration (12100 μM) and control group, the MDA level concentration-dependently decreased in the cells




Figure 2: The protective effects of hydro-alcoholic extract of *R. turkestanicum* against H_2O_2 -induced BAE-1 cells toxicity. The cells were pre-treated with different concentrations of *R. turkestanicum* for 24 h followed by 30 min incubation with H_2O_2 , and the cell viability was quantified by MTT assay. The data were expressed as percentage viability of control. The results are mean \pm SEM. The test was done in triplicate ($n = 3$). (***) $P < 0.001$ compared with H_2O_2 , (###) $P < 0.001$ compared with control




Figure 3: Effects of hydro-alcoholic extract of *R. turkestanicum* on ROS production in BAE-1 cells. The cells were pre-treated with different concentrations of *R. turkestanicum* and quercetin for 24 h followed by 30 min incubation with H₂O₂. The intracellular ROS was estimated via fluorescence intensity. The results were expressed as percentage of fluorescence intensity of control. The results are mean ± SEM. The test was done in triplicate (n = 3). (***P < 0.001 compared to H₂O₂, ###P < 0.001 compared with control)

treated with the extract. Similarly, significant differences were observed between the cells treated with the extract at 12100 µM and quercetin (P < 0.001).

Hydro-alcoholic extract of *R. turkestanicum* reduced apoptotic cells

As shown in Figure 5, H₂O₂ elevated apoptotic cells (P < 0.001) while quercetin and extract counteracted apoptotic cell following H₂O₂-induced apoptosis.

Discussion

In this research, we evaluated the protective effects of *R. turkestanicum* hydro-alcoholic extract against H₂O₂-induced toxicity in BAE-1 cells for the first time. H₂O₂ elevated cell death, apoptotic cells, ROS generation, and MDA at a dose of 200 µM. Pretreatment of cells with the extract reduced cell death, apoptotic cells, and oxidative stress via attenuation of lipid peroxidation and ROS production, dose-dependently. Quercetin as an active ingredient which is found in *R. turkestanicum*^[20] restored cell viability and decreased oxidative stress following H₂O₂-induced oxidative stress at a dose of 30.23 µg/ml. *R. turkestanicum* reduced doxorubicin-induced toxicity in cardiomyocytes (H9C2 cells),^[12] glutamate toxicity in PC12 cells.^[18] The protective effects of this herb against toxic agents such as gentamicin,^[14] mercuric chloride,^[15] hexachlorbutadien,^[16] and cisplatin^[13] have been also demonstrated. Another study showed that *R. turkestanicum* prevented cardiac injury following STZ-induced diabetic in rats via attenuation of lactate dehydrogenase and creatine phosphokinase.^[19] These studies have revealed the protective effects of *R. turkestanicum* may be related to the presence of active ingredients that scavenge free radicals




Figure 4: Effects of hydro-alcoholic extract of *R. turkestanicum* on lipid peroxidation in BAE-1 cells. The cells were pre-treated with different concentrations of *R. turkestanicum* and quercetin for 24 h followed by 30 min incubation with H₂O₂. The results are mean ± SEM. The MDA level was determined via TBARS fluorescence intensity. The results were expressed as percentage of fluorescence intensity of control. The test was done in triplicate (n = 3). (*P < 0.05, ***P < 0.001 compared to H₂O₂, ###P < 0.001 compared with control)

and attenuate oxidative stress.^[20] Bhushan *et al.* (2007) reported that *R. turkestanicum* induced apoptosis through ROS generation in human leukemic cells, HL60 and NB4. ROS formation dose-dependently has been elevated in the cells treated with *R. turkestanicum*. Therefore, *R. turkestanicum* and its components may act as both pro-oxidant and antioxidant, depending on the redox state of the biological environment.^[24] Rheum species is composed of different ingredients including anthrones, anthraquinones, quercetin, resveratrol, anthocyanins, acylglucosides, stilbenes, organic acids, and vitamins.^[20] The presence of anthraquinone derivatives including emodin, aloe-emodin, rhein, chrysophanol, physcion, and danthron as the main biologically active constituents of *Rheum* genus including *R. turkestanicum* have been identified.^[25,26] Anti-inflammatory and anti-apoptotic effects of emodin have reduced myocardial infarction in rat heart via elevation of antioxidant capacity.^[27] Moreover, it suppresses the expression of Toll-like receptor 4 and p38 mitogen activating protein kinase following viral myocarditis.^[28] Quercetin counteracted cardiac inflammation after ischemia-reperfusion by suppressing the activity of signal transducer and activator of transcription 3 (STAT3).^[29] Also, quercetin declined infarct size via activating the PI3K/Akt signaling pathway and modulating the expression of Bcl-2 and Bax proteins.^[30] Additionally, it reduced doxorubicin-induced cardiotoxicity in H9C2 cells and mice by the upregulation of Bmi-1 expression and suppression of oxidative stress.^[31] On the basis of the *in vitro* and *in vivo* studies, resveratrol revealed cardio-protective properties against doxorubicin toxicity through enhancing the antioxidant enzymes activity and attenuation of pro-apoptotic proteins such as




Figure 5: Effects of hydro-alcoholic extract of *R. turkestanicum* on BAE-1 apoptotic cells. The cells were pretreated with different concentrations of *R. turkestanicum* and quercetin for 24 h followed by 30 min with H_2O_2 . They were stained with PI for flow cytometric analysis. The flow cytometry histograms representing the cells with reduced DNA content accumulated in the sub-G1 region

p53, Bax, and caspase3.^[32-34] Resveratrol reduced ROS generation following palmitic acid-induced oxidative stress in human aortic endothelial cells.^[35] Rhein decreased H_2O_2 -induced toxicity in human umbilical vein endothelial cells via inhibition of ROS production and apoptosis.^[21] According to these findings, the protective effect of extract against oxidative stress may be mediated via active ingredients.

Conclusion

Our findings showed the protective effect of *R. turkestanicum* and quercetin against H_2O_2 -induced toxicity in endothelial cells. These beneficial properties are related to the presence of active ingredients, probably by suppression of free radicals, lipid peroxidation, and apoptotic cell death. The present study proposed that this extract can be a potential cardio-protective agent in the prevention of cardiovascular disease. However, more investigations are needed to warrant these findings.

Acknowledgments

This study financially supported by Vice Chancellor for Research and Technology (grant number 970640), Mashhad University of Medical Sciences, Mashhad, Iran.

Authors' contribution

Azar Hosseini, Mohammad Soukhtanloo, BizhanMalaekheh-Nikouei designed the study. Azar Hosseini, Arezoo Rajabian, Sahar Sheikh conducted the experiments. Azar Hosseini and Arezoo Rajabian wrote and revised the manuscript.

Financial support and sponsorship

This paper is a part of a Pharm.D thesis and was financially supported by Mashhad University of Medical Sciences.

Conflicts of interest

There are no conflicts of interest.

Received: 20 Oct 19 **Accepted:** 22 Apr 20

Published: 19 Aug 20

References

- Grover-Páez F, Zavalza-Gómez AB. Endothelial dysfunction and cardiovascular risk factors. *Diabetes Res Clin Pract* 2009;84:1-10.
- Martin BJ, Anderson TJ. Risk prediction in cardiovascular disease: The prognostic significance of endothelial dysfunction. *Can J Cardiol* 2009;25:15A-20A.
- Siti HN, Kamisah Y, Kamsiah J. The role of oxidative stress, antioxidants and vascular inflammation in cardiovascular disease (a review). *Vascul. Pharmacol* 2015;71:40-56.
- Sifuentes-Franco S, Padilla-Tejeda DE, Carrillo-Ibarra S, Miranda-Díaz AG. Oxidative stress, apoptosis, and mitochondrial function in Diabetic Nephropathy. *Int J Endocrinol* 2018;2018:1875870.
- Savini I, Catani MV, Evangelista D, Gasperi V, Avigliano L. Obesity-associated oxidative Stress: Strategies finalized to improve redox state. *Int J Mol Sci* 2013;14:10497-538.
- Cai H. Hydrogen peroxide regulation of endothelial function: Origins, mechanisms, and consequences. *Cardiovasc Res* 2005;68:26-36.
- Rahiman N, Akaberi M, Sahebkar A, Emami SA, Tayarani-Najaran Z. Protective effects of saffron and its active components against oxidative stress and apoptosis in endothelial cells. *Microvasc Res* 2018;118:82-9.
- Wongpradabchai S, Chularojmontri L, Phornchirasilp S,

- Wattanapitayakul SK. Protective effect of phyllanthus emblica fruit extract against hydrogen peroxide-induced endothelial cell death. *J Med Assoc Thai* 2013;96(Suppl 1):40-8.
9. Safaeian L, Sajjadi SE, Javanmard SH, Montazeri H, Samani F. Protective effect of Melissa officinalis extract against H₂O₂-induced oxidative stress in human vascularendothelial cells. *Res Pharm Sci* 2016;11:383-9.
 10. Bihanta M, Hosseini A, Ghorbani A, Boroushaki MT. Protective effect of pomegranate seed oil against H₂O₂-induced oxidative stress in cardiomyocytes. *Avicenna J Phytomed* 2017;7:46-53.
 11. Shiezadeh F, Mousavi SH, Amiri MS, Iranshahi M, Tayarani-Najaran Z, Karimi G. Cytotoxic and apoptotic potential of Rheum turkestanicum Janisch root extract on human cancer and normal cells. *Iran J Pharm Res* 2013;12:811-9.
 12. Hosseini A, Rajabian A. Protective effect of *Rheum turkestanicum* root against doxorubicin-induced toxicity in H9c2 cells. *Rev Bras Farmacogn* 2016;26:347-51.
 13. Hosseini A, Fanoudi S, Mollazadeh H, Aghaei A, Boroushaki MT. Protective effect of Rheum turkestanicum against cisplatin by reducing oxidative stress in kidney tissue. *J Pharm Bioallied Sci* 2018;10:66-71.
 14. Boroushaki MT, Fanoudi S, Mollazadeh H, Boroumand-Noughabi S, Hosseini A. Reno-protective effect of Rheum turkestanicum against gentamicin-induced nephrotoxicity. *Iran J Basic Med Sci* 2019;22:328-33.
 15. Hosseini A, Rajabian A, Fanoudi S, Farzadnia M, Boroushaki MT. Protective effect of *Rheum turkestanicum* root against mercuric chloride-induced hepatorenal toxicity in rats. *Avicenna J Phytomed* 2018;8:488-97.
 16. Boroushaki MT, Fanoudi S, Rajabian A, Boroumand S, Aghaei A, Hosseini A. Evaluation of *Rheum turkestanicum* in hexachlorobutadien-induced renal toxicity. *Drug Res (Stuttg)* 2019;69:434-8.
 17. Fei Y, Wang J, Peng B, Peng J, Hu JH, Zeng ZP. Phenolic constituents from Rheum nobile and their antioxidant activity. *Nat Prod Res* 2017;31:2842-9.
 18. Rajabian A, Sadeghnia HR, Moradzadeh M, Hosseini A. Rheum turkestanicum reduces glutamate toxicity in PC12 and N2a cell lines. *Folia Neuropathol* 2018;56:354-61.
 19. Hosseini A, Mollazadeh H, Amiri MS, Sadeghnia HR, Ghorbani A. Effects of a standardized extract of Rheum turkestanicum Janischew root on diabetic changes in the kidney, liver and heart of streptozotocin-induced diabetic rats. *Biomed Pharmacother* 2017;86:605-11.
 20. Singh P, Rawat M. Phytochemistry and biological activity perspectives of Rheum species. *Natural Products J* 2016;6:84-93.
 21. Zhong XF, Huang GD, Luo T, Deng ZY, Hu JN. Protective effect of rhein against oxidative stress-related endothelial cell injury. *Mol Med Rep* 2012;5:1261-16.
 22. Hosseini A, Shafiee-Nick R, Mousavi SH. Combination of *Nigella sativa* with Glycyrrhiza glabra and Zingiber officinale augments their protective effects on doxorubicin-induced toxicity in h9c2 cells. *Iran J Basic Med Sci* 2014;17:993-1000.
 23. Buege JA, Aust SD. Microsomal lipid peroxidation. *Methods Enzymol* 1978;53:302-10.
 24. Moradzadeh M, Rajabian A, Aghaei A, Hosseini A, Sadeghnia HR. *Rheum turkestanicum* induced apoptosis through ROS without a differential effect on human leukemic cells. *Jundishapur J Nat Pharm Prod* 2019;14:e12198.
 25. Dorsey J F, Kao GD. Aloe (-emodin) for cancer? More than just a comforting salve. *Cancer Biol Ther* 2007;6:89-90.
 26. Alam MM, Javed K and Jafri MA. Effect of Rheum emodi (Revand Hindi) on renal functions in rats. *J Ethnopharmacol* 2005;96:121-5.
 27. WuY, Tu X, Lin G, Xia H, Huang H, Wan J, et al. Emodin-mediated protection from acute myocardial infarction via inhibition of inflammation and apoptosis in local ischemic myocardium. *Life Sci* 2007;81:1332-8.
 28. Zhang Y, Lin C, Yang X, Wang Y, Fang Y, Wang F. Effect of emodin on the expression of TLR4 and P38MAPK in mouse cardiac tissues with viral myocarditis. *Int J Clin Exp Pathol* 2016;9:10839-45.
 29. Chen YW, Chou HC, Lin ST, Chen YH, Chang YJ, Chen L, et al. Cardioprotective effects of quercetin in cardiomyocyte under ischemia/reperfusion injury. *Evid Based Complement Alternat Med* 2013;2013:364519.
 30. Wang Y, Zhang Z, Wu Y, Ke J, He X, Wang Y. Quercetin postconditioning attenuates myocardial ischemia/reperfusion injury in rats through the PI3K/Akt pathway. *Braz J Med Biol Res* 2013;46:861-7.
 31. Dong Q, Chen L, Lu Q, Sharma S, Li L, Morimoto S, et al. Quercetin attenuates doxorubicin cardiotoxicity by modulating B mi-1 expression. *Br J Pharmacol* 2014;171:4440-54.
 32. Tatlidede E, Şehirli Ö, Velioglu-Öğünç A, Çetinel Ş, Yeğen BÇ, Yarat A, et al. Resveratrol treatment protects against doxorubicin-induced cardiotoxicity by alleviating oxidative damage. *Free Radic Res* 2009;43:195-205.
 33. Danz EDB, Skramsted J, Henry N, Bennett JA, Keller RS. Resveratrol prevents doxorubicin cardiotoxicity through mitochondrial stabilization and the Sirt1 pathway. *Free Radic Biol Med* 2009;46:1589-97.
 34. Sin TK, Tam BT, Yung BY, Yip SP, Chan LW, Wong CS, et al. Resveratrol protects against doxorubicin-induced cardiotoxicity in aged hearts through the SIRT1-USP7 axis. *J Physiol* 2015;593:1887-99.
 35. Jeong SO, Son Y, Lee JH, CheongYK, Park SH, Chung HT, et al. Resveratrol analog piceatannol restores the palmitic acid-induced impairment of insulin signaling and production of endothelial nitric oxide via activation of anti-inflammatory and antioxidative heme oxygenase-1 in human endothelial cells. *Mol Med Rep* 2015;12:937-44.