

## Role of Obesity in the Tumorigenesis of Gastric Cancer

### Abstract

Gastric cancer as a common cancer is a multi-factorial disease that is dependent on parallel effects of environment and genetics. Endogenous and host factors, including gender and several genetic backgrounds are known risk factors also many environmental factors, including smoking, diet, infection and increasing body weight and body mass index (BMI) are associated with the gastric cancer. Epidemiological data have consistently demonstrated a positive relation between obesity and gastric cancer, whereas mechanistic studies have sought to uncover obesity related carcinogenic pathways. Biological mechanisms and the relationship between obesity and cancer are complex and not well understood. Different effective factors include obesity-related hormones and adipokines, growth factors, modulation of energy balance and calorie restriction, inflammatory processes and multiple signaling pathways that affect cancer cell promotion and progression. In this review, we will discuss the recent advances in the understanding of the association of obesity changes in the gastric cancer.

**Keywords:** Adipokine, cytokine, gastric cancer, inflammation, obesity

### Introduction

Gastric cancer is the fifth most common cancer and the third leading cause of cancer death worldwide,<sup>[1]</sup> with a wide variation in incidence rates across different geographical areas.<sup>[2]</sup> Gastric cancer is a multi-factorial disease that is dependent to parallel effects of environment and genetics. Endogenous and host factors, including those related to male gender, and several genetic backgrounds are known risk factors to a lesser extent.<sup>[3]</sup>

Many environmental factors, including smoking, high salt intake and a diet with an insufficient level of antioxidants are involved in the pathogenesis of gastric cancer.<sup>[4]</sup> Several epidemiological studies have investigated the effects of increasing body weight and body mass index (BMI) on the risk of gastric cancer.<sup>[5,6]</sup>

Obesity is closely linked to an increase adipose tissue with dysfunction. The results of these changes are chronic inflammation and adverse interplay of sex steroids, endocrine hormones and various adipokines. Epidemiological data have consistently demonstrated a positive relation between obesity and gastric cancer, whereas

mechanistic studies have sought to uncover obesity related carcinogenic pathways.<sup>[7]</sup>

In this review, we will discuss the recent advances in the understanding of the obesity changes and association of chronic inflammation and adverse interplay of sex steroids, endocrine hormones and various adipokines with gastric cancer.

### Obesity and Cancer

Obesity is an important public health problem worldwide. Based on the World Health Organization (WHO) criteria, overweight and obesity are defined as abnormal or excessive fat accumulation that may impair health. Body mass index (BMI) is commonly used to classify overweight and obesity in adults. The WHO definition is a BMI greater than or equal to 25 is overweight and a BMI greater than or equal to 30 is obese.

According to the WHO report in 2014, more than 1.9 billion adults aged 18 years and older were overweight. The worldwide prevalence of obesity more than doubled between 1980 and 2014. Overall, 11% of men and 15% of women were obese.<sup>[8]</sup>

Epidemiological studies have demonstrated that obesity is associated with increased risk of several cancer types, including gastric, colon, endometrium, pancreas,

### Masoumeh Mohammadi

Cellular and Molecular  
Endocrine Research Center,  
Research Institute for Endocrine  
Sciences, Shahid Beheshti  
University of Medical Sciences,  
Tehran, Iran

#### Address for correspondence:

Dr. Masoumeh Mohammadi,  
Cellular and Molecular  
Endocrine Research Center,  
Research Institute for  
Endocrine Sciences, Shahid  
Beheshti University of Medical  
Sciences, Tehran, Iran.  
E-mail: [m-gorgian@yahoo.com](mailto:m-gorgian@yahoo.com)

#### Access this article online

**Website:**  
[www.ijpvmjournal.net/www.ijpvm.ir](http://www.ijpvmjournal.net/www.ijpvm.ir)

**DOI:**  
10.4103/ijpvm.IJPVM\_153\_19

#### Quick Response Code:



**How to cite this article:** Mohammadi M. Role of obesity in the tumorigenesis of gastric cancer. *Int J Prev Med* 2020;11:148.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: [reprints@medknow.com](mailto:reprints@medknow.com)

postmenopausal breast, esophagus, gallbladder, kidney, liver, and hematological malignancy.<sup>[9]</sup> Also, obesity is associated with an increase in cancer mortality. Among patients with a BMI  $\geq 40$  kg/m<sup>2</sup> compared with patients with a normal BMI, mortality from all causes of cancer was found to be 52% higher in men and 62% higher in women.<sup>[10]</sup>

Biological mechanisms and the relationship between obesity and cancer are complex and not well understood. Different effective factors include obesity-related hormones and adipokines, growth factors, modulation of energy balance and calorie restriction, inflammatory processes and multiple signaling pathways that, affect cancer cell promotion and progression.<sup>[6,11]</sup> Understanding of mechanisms which obesity may induce cancer development is important for prevention of cancer, also to improve outcomes for obese cancer patients.

Possible mechanisms linking obesity with gastric cancer may include obesity associated insulin resistance and an abnormally increased blood level of insulin like growth factor (IGF) and altered levels of adipokines such as adiponectin, leptin, resistin and visfatin.<sup>[12]</sup> It has been established that obesity is associated with higher levels of proinflammatory cytokines, including Tumor necrosis factor  $\alpha$  (TNF- $\alpha$ ), Interleukine-6 (IL-6) and C-reactive protein (CRP). Inflammation may stimulate cancer development by activation of Nuclear factor- $\kappa$ B (NF- $\kappa$ B) complex pathway.<sup>[11,13]</sup> Also, steroid hormones, including estrogen, androgens, progesterone, and adrenal steroids are associated with energy balance and progression of several types of cancer.<sup>[14]</sup>

### Factors involved in obesity associated gastric cancer and their mechanisms

#### Insulin and Insulin like growth factor (IGF)

Obesity is accompanied by increased insulin independently of type 2 diabetes, presumably due to increased need for metabolic processes energy.<sup>[15]</sup> According to the findings of several studies, insulin contributes to increased cancer risk. Insulin as a mitogenic hormone in muscle, liver and adipose tissue can increase tissue mass, augment glucose uptake, and finally synthesize nutrients. These anabolic effects are not directly related to carcinogenesis unclear, but could involve the activation of IGF system which is mitogenic under conditions of hyperglycemia or hyperinsulinemia, independently of type 2 diabetes, and alter levels of IGF binding protein-1 (IGFBP1) and -2 (IGFBP2) which accompanied by increased bioavailability of IGFs.<sup>[16]</sup>

In the obese patients, have been reported higher circulating levels of IGF-I than non-obese patients with hyperinsulinemia.<sup>[17]</sup> Insulin like growth factor-1 (IGF-1) is a mitogenic and anti-apoptotic factor, mainly produced by the liver, inhibits apoptosis and stimulates cell division therefore could contribute to the development of

cancer.<sup>[18]</sup> Insulin and IGF-1 through the phosphatidylinositol 3-kinase (PI3K)/Akt target of rapamycin (mTOR) pathway and Ras-raf-MAPK signaling which alter the expression of genes involved in cell growth and proliferation<sup>[19,20]</sup> Figure 1. In addition, IGFs and insulin like growth factor-1 receptors (IGF-1R) and insulin are over expressed in various cancer tissues.<sup>[21]</sup> IGF-1Rs have tyrosine kinase activity and after binding of their ligands, these receptors activate and promote cell migration and redistribution of a- and b-catenins and E-cadherin from adherent junctions into the cytoplasm and promote tumorigenesis.<sup>[16,22]</sup>

Epidemiological data have consistently demonstrated a positive relation between type 2 diabetes mellitus and gastric cancer. The molecular basis for this association is unclear but could involve IGF-I signaling, which is mitogenic under conditions of hyperglycemia or hyperinsulinemia.<sup>[23]</sup>

#### Adipokines

The adipose tissue is a complex endocrine organ. In obesity, adipose tissues have a state of low grade chronic inflammation and presence inflammatory cells (e.g. macrophages and lymphocytes). In fact, adipose tissue involved in carcinogenesis via dysregulated secretion of various adipokines, which are produced by white adipose tissue.<sup>[24,25]</sup> These adipokines have been implicated in cancer development through effects on various inflammatory pathways, insulin resistance and lipolysis.<sup>[26]</sup>

More than 15 adipokines have been reported in the literatures to be associated with cancers.<sup>[27]</sup> Among these




Figure 1: IGF associated gastric cancer and its mechanisms: By binding with IGFs and IGF receptors activate signaling pathways such as RAS/MAPK and PI-3K/AKT, thus resulting in the functions of proliferation, survival, apoptosis and differentiation in cancer cells

adipokines some adipokines such as adiponectin are protective against tumorigenesis and its serum levels are usually decreased in the patients with cancer while the circulating levels of majority of pro-inflammatory adipokine levels, such as leptin, resistin, and visfatin increase in cancers.<sup>[24]</sup> In continues, refers to a series of these adipokines that act as circulating hormones in gastric.

### Resistin

Resistin is a member of family of cysteine-rich proteins called "Resistin-like molecules" (RELMs). Its gene, called Retn mapping to the p13.3 band of chromosome,<sup>[28]</sup> encoded a 12.5-kDa protein.<sup>[29]</sup> It is expressed at very low levels, in human adipose cells, whereas high levels of resistin are expressed in mononuclear leukocytes, macrophages, spleen, and bone marrow cells.<sup>[30,31]</sup>

Resistin has identified as an important mediator in inflammation associated cancers. Few studies, in human, have been investigating the association of resistin and RELMs with gastric cancer. Zheng *et al.* reported a higher expression of RELM $\beta$  in intestinal type compared to diffuse type gastric carcinomas. In addition, RELM $\beta$  correlated positively with tumor differentiation and inversely with tumor infiltration, lymph node metastasis and heparanase expression.<sup>[32]</sup> About resistin in gastric cancer, Nakajima *et al.* showed that resistin correlate significantly with stage progression and suggested its use as a biomarker for gastric cancer progression.<sup>[12]</sup> In a previous study, we showed that resistin doesn't express by gastric cancer cell and exogenously by paracrine pathway causes gastric cancer progression.<sup>[33]</sup> In another study, we showed that exogenous resistin induces gastric cancer cell proliferation by increases human telomerase reverse transcriptase (hTERT) gene expression.<sup>[34]</sup>

Hsieh *et al.* showed that resistin induced stromal cell-derived factor-1 (SDF-1) upregulation by activation of Toll-like receptor 4 (TLR4), P38 mitogen-activated protein kinases (p38 MARK) and nuclear factor NF- $\kappa$ B. SDF-1 is involved in the carcinogenesis of human gastric cancer, where it stimulates angiogenesis and favors metastasis of tumor cells to distant organs.<sup>[35]</sup>

### Visfatin

Visfatin, a novel adipocyte derived adipokine, is highly enriched in the visceral fat and its plasma levels increase during the development of obesity.<sup>[36]</sup> Visfatin is also known as pre-B cell enhancing factor (PBEF), a growth factor for early B cell proliferation,<sup>[37]</sup> and it is the secretory form of nicotinamide phosphoribosyl-transferase (Nampt),<sup>[38]</sup> the rate-limiting enzyme of mammalian nicotinamide adenine dinucleotide (NAD) biosynthesis.

As yet visfatin receptor has not been identified but in studies on diabetes had indicated that visfatin can significantly regulate insulin secretion. Presumably visfatin with use of insulin receptor and intracellular signaling can

regulate insulin secretion from pancreatic beta cells and affects a variety of metabolic and stress responses.<sup>[39,40]</sup>

Visfatin have three major functions: Growth Factor, cytokine and nicotinamide phosphoribosyl transferase, therefore increasing of visfatin have several effects. Recent studies have shown that over-expression of visfatin is important in the carcinogenesis in several types of cancers.<sup>[41,42]</sup> several studies have shown visfatin is over-expressed in human gastric cancer tissues and established gastric cancer cells<sup>[33,43]</sup> also reported that high plasma visfatin levels gradually increase with stage progression of gastric cancer.<sup>[12]</sup>

Visfatin by up regulation of cyclin D1 and cdk2 expression activates G1-S phase cell cycle progression. Visfatin also increases the expression of matrix metalloproteinases 2, matrix metalloproteinases 9, and vascular endothelial growth factor genes, therefore induces metastasis and angiogenesis of cancer.<sup>[41,44]</sup> In several studies have shown, visfatin can regulate a variety of different signaling pathways such as Signal transducer and activator of transcription 3 (STAT3), PI3K/Akt and extracellular signal-regulated protein kinases 1 and 2 (ERK1/2).<sup>[45,46]</sup>

These findings about visfatin show its role in carcinogenesis and cancer progression. Few studies, in human, have been investigated the association of visfatin with gastric cancer, in a previous study, we showed that visfatin expresses by gastric cancer cell and endogenously causes gastric cancer progression.<sup>[33]</sup> In another study, we showed that visfatin upregulation of human telomerase reverse transcriptase (hTERT) gene expression can cause gastric cancer progression.<sup>[47]</sup>

### Leptin

Leptin gene is located on chromosome 7 in humans and encodes a 16 kDa protein with 167 amino acids.<sup>[48]</sup> Leptin is produced primarily in the adipocytes of white and brown adipose tissue, placenta, ovaries, bone marrow, skeletal muscle, mammary epithelial cells, pituitary, liver, gastric chief cells and P/D1 cells.<sup>[49,50]</sup> In obese individuals, there is a higher circulating concentration of leptin than normal weight individuals. Leptin reduces appetite as a circulating signal, but obese individuals show resistance to leptin.<sup>[51]</sup>

Leptin stimulates the proliferation, migration, and invasion of tumor cells. Several studies have investigated the expression of the leptin in normal and malignant gastric tissue<sup>[52-60]</sup> and have shown there is a positive correlation between expression of leptin and its receptor.<sup>[52]</sup> Also, high leptin and its receptor expression were associated with aggressiveness of tumor.<sup>[52,57,58]</sup> In fact, it is a key player in the obesity and cancer link.

*In vitro* studies have shown leptin causes increased cell proliferation in gastric carcinoma cell lines AGS,<sup>[54]</sup> MKN-28,<sup>[55,56]</sup> MKN-74<sup>[55]</sup> and HS746T,<sup>[56]</sup> via activating a

number of signaling pathways, JAK2/STAT3 and STAT5, IRS/PI3K, SHP2/MAPK, and AMPK/ACC.<sup>[55,56,61-63]</sup>

### Adiponectin

Adiponectin is an adipose tissue derived hormone, which is down regulated in obesity. Circulating concentrations of adiponectin are 3 to 30 ng/mL that is expressed in differentiated adipocytes.<sup>[64,65]</sup>

Adiponectin has an inverse association with total body fat mass and displays, protective actions on the development of various obesity related diseases such as cancer.<sup>[66]</sup> Epidemiologic studies show that low level of adiponectin has an inverse association with cancer progression as well as advanced disease.<sup>[67,68]</sup> Lower plasma adiponectin levels have also been found in patients with gastric cancer, especially in upper gastric cancer, compared with healthy control subjects, and were inversely correlated with tumor size, depth of invasion, and metastasis stage (76). This inverse correlation is due to anti-inflammatory, antitumor, and antiangiogenic effects of adiponectin (75). The anti-inflammatory effects of adiponectin are supported by an observed inverse correlation between plasma levels of adiponectin and C-reactive protein.<sup>[69]</sup> The antitumor and anti-angiogenic effects of adiponectin are supported by an observed induction of apoptosis and inhibits the proliferation of gastric cancer cell lines through adiponectin receptors AdipoR1 and AdipoR2 (78).

### Inflammatory cytokines

Obesity is associated with chronic low-grade inflammation (or meta-inflammation), characterized by increased inflammatory signaling, abnormal cytokine production and immune activation. The connection between inflammation and cancer has been studied for years, and chronic inflammation is thought to be a key contributor to tumor development.<sup>[70]</sup>

Inflammatory cytokines and chemokines, including TNF- $\alpha$ , IL-6, and CRP are more deregulated in the gastric cancer. In continues, refers to a series of these inflammatory cytokines that act as contributor to tumor development.

### Tumor necrosis factor $\alpha$

Tumor necrosis factor  $\alpha$  is a 26-kDa transmembrane protein and a powerful inflammatory cytokine. TNF- $\alpha$ , first identified as an antitumor agent, but now also is recognized as a tumor promoting cytokine that links inflammation and cancer.<sup>[71]</sup> Recent studies have indicated that TNF-alpha overexpression in adipose tissue is a common feature in human and animal models of obesity and plays an important role in mediating the insulin resistance by over-expression in fat tissue.<sup>[72,73]</sup>

In fact, TNF- $\alpha$  contributes to the deregulation of the insulin signaling pathway, including serine phosphorylation of insulin-receptor substrate (IRS) proteins by c-jun

N terminal kinase (JNK) and an inhibitor of nuclear factor- $\kappa$ B kinase (IKK)<sup>[74,75]</sup> and induce inhibitory serine 307 (Ser307) phosphorylation of IRS-1 which decreases activation of insulin mediated Akt/PI3K/mTOR pathway.<sup>[76,77]</sup> Zhao *et al.* investigated link between over-expression of CXCR4 and an obvious involvement in gastric cancer metastasis. They showed that *H. pylori* upregulates CXCR4 expression in gastric cancer through TNF-alpha.<sup>[78]</sup> Oshima *et al.* investigated the role of TNF- $\alpha$  in gastric tumorigenesis and indicated that TNF- $\alpha$  signaling through Tumor necrosis factor receptor 1 (TNFR1) is important for gastric tumor development. They suggested that the activation of TNF- $\alpha$ /TNFR1 signaling in the tumor promotes gastric cancer and development through induction of Nicotinamide adenine dinucleotide phosphate oxidase organizer 1 (Noxo1) and G Protein Subunit Alpha 14 (Gna14), contribute to maintaining the tumor cells in an undifferentiated state.<sup>[79]</sup>

### Interleukin 6

IL-6 is produced by a variety of different cell types such as fibroblasts, endothelial cells, macrophages, T cells and myocytes. In obesity, adipose tissue, increase in adipocyte size, number and fat mass that is associated with recruitment and activation of T lymphocytes and macrophages in this tissue with the production of IL-6.<sup>[80,81]</sup>

The principal signaling mechanism of IL-6 involves activation of the transducer and activator of transcription 3 (STAT3) transcription factor and subsequent regulation of gene transcription. Other signaling pathways activated by IL-6 include PI3K and AMPK signaling pathways.

It has been suggested that IL-6 play a potential role in the pathogenesis and growth of malignancies. Production of IL-6 by gastric cancer cell lines has been reported and act in a paracrine and autocrine manner in tumor cells. It has also been shown that IL-6 levels correlated with clinicopathological features of gastric cancer.<sup>[82,83]</sup>

### C-reactive protein (CRP)

C-reactive protein is a marker of systemic inflammation produced mainly by the liver under the stimulation of adipocyte derived proinflammatory cytokines and elevated in obesity.<sup>[84]</sup> Serum CRP elevates in many malignancies, and it is a prognostic indicator of malignant potential. However, the prognostic value of CRP in gastric cancer remains uncertain, but in many studies have been shown that the serum CRP level is abnormally elevated in gastric cancer patients and might be a potential prognostic factor.<sup>[85]</sup> Recently, Chen *et al.* evaluated the significance of serum CRP as a biomarker of long term survival in patients with advanced gastric cancer. They showed that elevated serum CRP baseline levels before chemotherapy were associated with reduced survival in patients with advanced gastric cancer. The rate of CRP



declining was also associated with survival. The CRP baseline concentration before and after chemotherapy may be used as predictive markers of advanced gastric cancer.<sup>[86]</sup>

### Sex hormones

Epidemiologic studies suggest differences in complications from obesity in men vs women, which determines differences in plasma levels of sex hormones.<sup>[65]</sup> Endogenous sex hormones are active in tumor cell growth and directly mediate the effect of obesity on cancer.

Data from studies have been shown that gastric cancer for most populations in both high and low incidence regions, in men is approximately double that of women. These differences cannot be explained by variations in environmental factors or *Helicobacter pylori* (H. pylori) infection, therefore female sex hormones have been proposed to be protective.

Constanza Camargo *et al.* performed a meta-analysis to examine associations of gastric cancer risk in women with sex hormone. Their analysis showed that longer exposure to estrogen effects of either ovarian or exogenous origin may decrease risk of gastric cancer.<sup>[1]</sup>

There is evidence that estrogens regulate the insulin receptor substrate-1 (IRS-1) expression and induce free radical mediated DNA damage and gene mutations in cells.<sup>[87]</sup> Also, has been shown that estrogen contribute to the decreasing insulin sensitivity, regulation of body adiposity and increasing leptin signaling pathways.<sup>[88]</sup> In women, leptin levels during the menstrual cycle correlate directly with the estrogen levels. Regulation of leptin levels with estrogen in women may play a protective role, although a high level of leptin is considered to be a risk factor for males to cancer development.<sup>[89]</sup> Overall, estrogen may prevent gastric cancer, but additional studies are warranted.

### Genetic factors

In recent years, several new genes have been discovered to be associated with gastric cancer. For example, Immune response-related genes, Gastric mucosal protective genes, Metabolic enzyme genes, DNA repair genes, Tumor suppressor genes, transforming growth factor b (TGF-b), and Vascular endothelial growth-factor (VEGF).<sup>[90]</sup> According to the epidemiological studies, about 65-80% of the variation in BMI is determined by genetic factors.<sup>[91]</sup> Obesity is associated with a risk of at least 20 different cancers.<sup>[92]</sup>

Genome-wide association studies in colorectal cancer have described altered macrophage-enriched metabolic network genes.<sup>[93]</sup> The single-nucleotide polymorphism associated with obesity is the fat mass and obesity related (FTO) gene, which may function through nutrient sensing but with no relation to gastrointestinal cancers.<sup>[65,94]</sup>

## Discussion

In this review, we attempted to summarize and discuss available data in order to better understand the association of obesity changes such as chronic inflammation and adverse interplay of sex steroids, endocrine hormones and various adipokines with the gastric cancers.

Several epidemiological studies have investigated the effect of increasing body weight and obesity on the risk of gastric cancer. Obesity is an inflammatory disease that leads to the onset of metabolic syndrome and cancer. Possible mechanisms linking obesity with gastric cancer may include obesity associated insulin resistance and an abnormally increased blood level of insulin like growth factor (IGF) and altered levels of adipokines such as adiponectin, leptin, resistin and visfatin.<sup>[12]</sup> It has been established that obesity is associated with higher levels of pro-inflammatory cytokines, including TNF- $\alpha$ , IL-6 and CRP are infiltrating the inflamed tissues and trigger pathways of activation of NF- $\kappa$ B complex pathway.<sup>[11,13]</sup> On the other hand, changes in sex steroids, including estrogen, androgens, progesterone, and adrenal steroids are associated with energy balance and progression of several types of cancer.<sup>[14]</sup>

In conclusion, characterization of the relationship between obesity and gastric cancers could help in the establishment of new preventive strategies and also in identification of new specific targets for cancer therapy, for improving life quality of patients and the quality of health services.

### Acknowledgments

This study was supported by Cellular and Molecular Endocrine Research Center, Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences. We are grateful to all individuals who participated in this project.

### Financial support and sponsorship

Research Institute for Endocrine Sciences, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

### Conflicts of interest

There are no conflicts of interest.

**Received:** 27 Apr 19 **Accepted:** 01 Aug 19

**Published:** 10 Sep 20

### References

1. Camargo MC, Goto Y, Zabaleta J, Morgan DR, Correa P, Rabkin CS. Sex hormones, hormonal interventions, and gastric cancer risk: A meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2012;21:20-38.
2. Parkin DM. International variation. *Oncogene* 2004;23:6329-40.
3. Guilford P, Hopkins J, Harraway J, Mcleod M, Mcleod N, Harawira P, *et al.* E-cadherin germline mutations in familial gastric cancer. *Nature* 1998;392:402-5.
4. Malekzadeh R, Derakhshan MH, Malekzadeh Z. Gastric cancer in Iran: Epidemiology and risk factors. *Arch Iran Med* 2009;12:576-83.

5. Gunter MJ, Leitzmann MF. Obesity and colorectal cancer: Epidemiology, mechanisms and candidate genes. *J Nutr Biochem* 2006;17:145-56.
6. Calle EE, Kaaks R. Overweight, obesity and cancer: Epidemiological evidence and proposed mechanisms. *Nat Rev Cancer* 2004;4:579-91.
7. Yang P, Zhou Y, Chen B, Wan HW, Jia GQ, Bai HL, *et al.* Overweight, obesity and gastric cancer risk: Results from a meta-analysis of cohort studies. *Eur J Cancer* 2009;45:2867-73.
8. Available from: <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>.
9. Lichtman MA. Obesity and the risk for a hematological malignancy: Leukemia, lymphoma, or myeloma. *Oncologist* 2010;15:1083-101.
10. Pi-Sunyer X. The medical risks of obesity. *Postgrad Med* 2009;121:21-33.
11. Harvey AE, Lashinger LM, Hursting SD. The growing challenge of obesity and cancer: An inflammatory issue. *Ann N Y Acad Sci* 2011;1229:45-52.
12. Nakajima TE, Yamada Y, Hamano T, Furuta K, Gotoda T, Katai H, *et al.* Adipocytokine levels in gastric cancer patients: Resistin and visfatin as biomarkers of gastric cancer. *J Gastroenterol* 2009;44:685-90.
13. Hursting SD, Berger NA. Energy balance, host-related factors, and cancer progression. *J Clin Oncol* 2010;28:4058-65.
14. Hursting SD, Lashinger LM, Wheatley KW, Rogers CJ, Colbert LH, Nunez NP, *et al.* Reducing the weight of cancer: Mechanistic targets for breaking the obesity-carcinogenesis link. *Best Pract Res Clin Endocrinol Metab* 2008;22:659-69.
15. Lakka HM, Salonen JT, Tuomilehto J, Kaplan GA, Lakka TA. Obesity and weight gain are associated with increased incidence of hyperinsulinemia in non-diabetic men. *Horm Metab Res* 2002;34:492-8.
16. Khan S, Shukla S, Sinha S, Meeran SM. Role of adipokines and cytokines in obesity-associated breast cancer: Therapeutic targets. *Cytokine Growth Factor Rev* 2013;24:503-13.
17. Brick DJ, Gerweck AV, Meenaghan E, Lawson EA, Misra M, Fazeli P, *et al.* Determinants of IGF1 and GH across the weight spectrum: From anorexia nervosa to obesity. *Eur J Endocrinol* 2010;163:185-91.
18. Finkelstein JW, Kream J, Ludan A, Hellman L. Sulfation factor (somatomedin): An explanation for continued growth in the absence of immunoassayable growth hormone in patients with hypothalamic tumors. *J Clin Endocrinol Metab* 1972;35:13-7.
19. Renehan AG, Frystyk J, Flyvbjerg A. Obesity and cancer risk: The role of the insulin-IGF axis. *Trends Endocrinol Metab* 2006;17:328-36.
20. Leroith D, Roberts CT Jr. The insulin-like growth factor system and cancer. *Cancer Lett* 2003;195:127-37.
21. Law JH, Habibi G, Hu K, Masoudi H, Wang MY, Stratford AL, *et al.* Phosphorylated insulin-like growth factor-1/insulin receptor is present in all breast cancer subtypes and is related to poor survival. *Cancer Res* 2008;68:10238-46.
22. Chan BT, Lee AV. Insulin receptor substrates (IRSs) and breast tumorigenesis. *J Mammary Gland Biol Neoplasia* 2008;13:415-22.
23. Shimoyama S. Diabetes mellitus carries a risk of gastric cancer: A meta-analysis. *World J Gastroenterol* 2013;19:6902-910.
24. Raucci R, Rusolo F, Sharma A, Colonna G, Castello G, Costantini S. Functional and structural features of adipokine family. *Cytokine* 2013;61:1-14.
25. Harvey AE, Lashinger LM, Hursting SD. The growing challenge of obesity and cancer: An inflammatory issue. *Ann N Y Acad Sci* 2011;1229:45-52.
26. Pérez-Hernández AI, Catalán V, Gómez-Ambrosi J, Rodríguez A, Frühbeck G. Mechanisms linking excess adiposity and carcinogenesis promotion. *Front Endocrinol (Lausanne)* 2014;5:65.
27. Arner E, Forrest AR, Ehrlund A, Mejhert N, Itoh M, Kawaji H, *et al.* Ceruloplasmin is a novel adipokine which is overexpressed in adipose tissue of obese subjects and in obesity-associated cancer cells. *PLoS One* 2014;9:e80274.
28. Urbanek M, Du Y, Silander K, Collins FS, Stepan CM, Strauss JF 3<sup>rd</sup>, *et al.* Variation in resistin gene promoter not associated with polycystic ovary syndrome. *Diabetes* 2003;52:214-7.
29. Meier U, Gressner AM. Endocrine regulation of energy metabolism: Review of pathobiochemical and clinical chemical aspects of leptin, ghrelin, adiponectin, and resistin. *Clin Chem* 2004;50:1511-25.
30. Lu SC, Shieh WY, Chen CY, Hsu SC, Chen HL. Lipopolysaccharide increases resistin gene expression *in vivo* and *in vitro*. *FEBS Lett* 2002;530:158-62.
31. Patel L, Buckels AC, Kinghorn IJ, Murdock PR, Holbrook JD, Plumpton C, *et al.* Resistin is expressed in human macrophages and directly regulated by PPAR gamma activators. *Biochem Biophys Res Commun* 2003;300:472-6.
32. Zheng LD, Tong QS, Weng MX, He J, Lv Q, Pu JR, *et al.* Enhanced expression of resistin-like molecule beta in human colon cancer and its clinical significance. *Dig Dis Sci* 2009;54:274-81.
33. Gorgian Mohammadi M, Hedayati M, Zarghami N, Ghaemmaghami S, Mohaddes M. Adipocyte derived hormones gene expression, resistin and visfatin, in AGS gastric cancer cell line. *Iran J Cancer Prev* 2013;6:165-9.
34. Mohammadi M, Hedayati M, Zarghami N, Ghaemmaghami S. Resistin effects on telomerase gene expression in AGS gastric cancer cell line. *Acta Endocrinol (Buchar)* 2016;12:145-9.
35. Hsieh YY, Shen CH, Huang WS, Chin CC, Kuo YH, Hsieh MC, *et al.* Resistin-induced stromal cell-derived factor-1 expression through toll-like receptor 4 and activation of p38 MAPK/NFκB signaling pathway in gastric cancer cells. *J Biomed Sci* 2014;21:59.
36. Fukuhara A, Matsuda M, Nishizawa M, Segawa K, Tanaka M, Kishimoto K, *et al.* Visfatin: A protein secreted by visceral fat that mimics the effects of insulin. *Science* 2005;307:426-30.
37. Samal B, Sun Y, Stearns G, Xie C, Suggs S, Mcniece I. Cloning and characterization of the cDNA encoding a novel human pre-B-cell colony-enhancing factor. *Mol Cell Biol* 1994;14:1431-7.
38. Imai S. Nicotinamide phosphoribosyltransferase (Nampt): A link between NAD biology, metabolism, and diseases. *Curr Pharm Des* 2009;15:20-8.
39. Brown JE, Onyango DJ, Ramanjaneya M, Conner AC, Patel ST, Dunmore SJ, *et al.* Visfatin regulates insulin secretion, insulin receptor signalling and mRNA expression of diabetes-related genes in mouse pancreatic beta-cells. *J Mol Endocrinol* 2010;44:171-8.
40. Revollo JR, Korner A, Mills KF, Satoh A, Wang T, Garten A, *et al.* Nampt/PBEF/Visfatin regulates insulin secretion in beta cells as a systemic NAD biosynthetic enzyme. *Cell Metab* 2007;6:363-75.
41. Kim JG, Kim EO, Jeong BR, Min YJ, Park JW, Kim ES, *et al.* Visfatin stimulates proliferation of MCF-7 human breast cancer cells. *Mol Cells* 2010;30:341-5.
42. Patel ST, Mistry T, Brown JE, Digby JE, Adya R, Desai KM, *et al.* A novel role for the adipokine visfatin/pre-B cell colony-enhancing factor 1 in prostate carcinogenesis. *Peptides* 2010;31:51-7.

43. Bi TQ, Che XM. Namp1/PBEF/visfatin and cancer. *Cancer Biol Ther* 2010;10:119-25.
44. Adya R, Tan BK, Punn A, Chen J, Randeve HS. Visfatin induces human endothelial VEGF and MMP-2/9 production via MAPK and PI3K/Akt signalling pathways: Novel insights into visfatin-induced angiogenesis. *Cardiovasc Res* 2008;78:356-65.
45. Adya R, Tan BK, Punn A, Chen J, Randeve HS. Visfatin induces human endothelial VEGF and MMP-2/9 production via MAPK and PI3K/Akt signalling pathways: Novel insights into visfatin-induced angiogenesis. *Cardiovasc Res* 2008;78:356-65.
46. Kim JY, Bae YH, Bae MK, Kim SR, Park HJ, Wee HJ, *et al.* Visfatin through STAT3 activation enhances IL-6 expression that promotes endothelial angiogenesis. *Biochim Biophys Acta* 2009;1793:1759-67.
47. Mohammadi M, Zarghami N, Hedayati M, Ghaemmaghami S, Yamchi RM, Mohaddes M. Visfatin effects on telomerase gene expression in AGS gastric cancer cell line. *Indian J Cancer* 2015;52:32-5.
48. Green ED, Maffei M, Braden VV, Proenca R, Desilva U, Zhang Y, *et al.* The human obese (OB) gene: RNA expression pattern and mapping on the physical, cytogenetic, and genetic maps of chromosome 7. *Genome Res* 1995;5:5-12.
49. Bado A, Levasseur S, Attoub S, Kermorgant S, Laigneau JP, Bortoluzzi MN, *et al.* The stomach is a source of leptin. *Nature* 1998;394:790-3.
50. Margetic S, Gazzola C, Pegg GG, Hill RA. Leptin: A review of its peripheral actions and interactions. *Int J Obes Relat Metab Disord* 2002;26:1407-33.
51. Considine RV, Sinha MK, Heiman ML, Kriauciunas A, Stephens TW, Nyce MR, *et al.* Serum immunoreactive-leptin concentrations in normal-weight and obese humans. *N Engl J Med* 1996;334:292-5.
52. Ishikawa M, Kitayama J, Nagawa H. Expression pattern of leptin and leptin receptor (OB-R) in human gastric cancer. *World J Gastroenterol* 2006;12:5517-22.
53. Zhao L, Shen ZX, Luo HS, Shen L. Possible involvement of leptin and leptin receptor in developing gastric adenocarcinoma. *World J Gastroenterol* 2005;11:7666-70.
54. Schneider R, Bornstein SR, Chrousos GP, Boxberger S, Ehninger G, Breidert M. Leptin mediates a proliferative response in human gastric mucosa cells with functional receptor. *Horm Metab Res* 2001;33:1-6.
55. Shida D, Kitayama J, Mori K, Watanabe T, Nagawa H. Transactivation of epidermal growth factor receptor is involved in leptin-induced activation of janus-activated kinase 2 and extracellular signal-regulated kinase 1/2 in human gastric cancer cells. *Cancer Res* 2005;65:9159-63.
56. Pai R, Lin C, Tran T, Tarnawski A. Leptin activates STAT and ERK2 pathways and induces gastric cancer cell proliferation. *Biochem Biophys Res Commun* 2005;331:984-92.
57. Zhao X, Huang K, Zhu Z, Chen S, Hu R. Correlation between expression of leptin and clinicopathological features and prognosis in patients with gastric cancer. *J Gastroenterol Hepatol* 2007;22:1317-21.
58. Hong SJ, Kwon KW, Kim SG, Ko BM, Ryu CB, Kim YS, *et al.* Variation in expression of gastric leptin according to differentiation and growth pattern in gastric adenocarcinoma. *Cytokine* 2006;33:66-71.
59. Sobhani I, Bado A, Vissuzaine C, Buyse M, Kermorgant S, Laigneau JP, *et al.* Leptin secretion and leptin receptor in the human stomach. *Gut* 2000;47:178-83.
60. Bolukbas FF, Kilic H, Bolukbas C, Gumus M, Horoz M, Turhal NS, *et al.* Serum leptin concentration and advanced gastrointestinal cancers: A case controlled study. *BMC Cancer* 2004;4:29.
61. Park HK, Ahima RS. Leptin signaling. *F1000Prime Rep* 2014;6:73.
62. Dalamaga M, Chou SH, Shields K, Papageorgiou P, Polyzos SA, Mantzoros CS. Leptin at the intersection of neuroendocrinology and metabolism: Current evidence and therapeutic perspectives. *Cell Metab* 2013;18:29-42.
63. Moon HS, Dalamaga M, Kim SY, Polyzos SA, Hamnvik OP, Magkos F, *et al.* Leptin's role in lipodystrophic and nonlipodystrophic insulin-resistant and diabetic individuals. *Endocr Rev* 2013;34:377-412.
64. Motoshima H, Wu X, Sinha MK, Hardy VE, Rosato EL, Barbot DJ, *et al.* Differential regulation of adiponectin secretion from cultured human omental and subcutaneous adipocytes: Effects of insulin and rosiglitazone. *J Clin Endocrinol Metab* 2002;87:5662-7.
65. Alemán JO, Eusebi LH, Ricciardiello L, Patidar K, Sanyal AJ, Holt PR. Mechanisms of obesity-induced gastrointestinal neoplasia. *Gastroenterology* 2014;146:357-73.
66. Ouchi N, Walsh K. Adiponectin as an anti-inflammatory factor. *Clin Chim Acta* 2007;380:24-30.
67. Barb D, Williams CJ, Neuwirth AK, Mantzoros CS. Adiponectin in relation to malignancies: A review of existing basic research and clinical evidence. *Am J Clin Nutr* 2007;86:858S-66S.
68. Wei T, Ye P, Peng X, Wu LL, Yu GY. Circulating adiponectin levels in various malignancies: An updated meta-analysis of 107 studies. *Oncotarget* 2016;7:48671-91.
69. Ouchi N, Kihara S, Funahashi T, Matsuzawa Y, Walsh K. Obesity, adiponectin and vascular inflammatory disease. *Curr Opin Lipidol* 2003;14:561-6.
70. Coussens LM, Werb Z. Inflammation and cancer. *Nature* 2002;420:860-7.
71. Balkwill F. Tumour necrosis factor and cancer. *Nat Rev Cancer* 2009;9:361-71.
72. Araújo EP, De Souza CT, Ueno M, Cintra DE, Bertolo MB, Carnevali JB, *et al.* Infliximab restores glucose homeostasis in an animal model of diet-induced obesity and diabetes. *Endocrinology* 2007;148:5991-7.
73. Hotamisligil GS, Peraldi P, Budavari A, Ellis R. IRS-1-mediated inhibition of insulin receptor tyrosine kinase activity in TNF- $\alpha$ -and obesity-induced insulin resistance. *Science* 1996;271:665.
74. Ghosh S, Karin M. Missing pieces in the NF- $\kappa$ B puzzle. *Cell* 2002;109:S81-96.
75. Lee YH, Giraud J, Davis RJ, White MF. c-Jun N-terminal kinase (JNK) mediates feedback inhibition of the insulin signaling cascade. *J Biol Chem* 2003;278:2896-902.
76. Aguirre V, Werner ED, Giraud J, Lee YH, Shoelson SE, White MF. Phosphorylation of Ser307 in insulin receptor substrate-1 blocks interactions with the insulin receptor and inhibits insulin action. *J Biol Chem* 2002;277:1531-7.
77. Gao Z, Hwang D, Bataille F, Lefevre M, York D, Quon MJ, *et al.* Serine phosphorylation of insulin receptor substrate 1 by inhibitor  $\kappa$ B kinase complex. *J Biol Chem* 2002;277:48115-21.
78. Zhao C, Lu X, Bu X, Zhang N, Wang W. Involvement of tumor necrosis factor- $\alpha$  in the upregulation of CXCR4 expression in gastric cancer induced by helicobacter pylori. *BMC Cancer* 2010;10:419.
79. Oshima H, Ishikawa T, Yoshida GJ, Naoi K, Maeda Y, Naka K, *et al.* TNF- $\alpha$ /TNFR1 signaling promotes gastric tumorigenesis through induction of Nox1 and Gna14 in tumor cells. *Oncogene* 2014;33:3820-9.

80. Ghigliotti G, Barisione C, Garibaldi S, Fabbi P, Brunelli C, Spallarossa P, *et al.* Adipose tissue immune response: Novel triggers and consequences for chronic inflammatory conditions. *Inflammation* 2014;37:1337-53.
81. Vozarova B, Weyer C, Hanson K, Tataranni PA, Bogardus C, Pratley RE. Circulating interleukin-6 in relation to adiposity, insulin action, and insulin secretion. *Obes Res* 2001;9:414-7.
82. Lukaszewicz-Zajac M, Mroczko B, Szmitkowski M. [The role of interleukin-6 and C-reactive protein in gastric cancer]. *Pol Merkur Lekarski* 2010;29:382-6.
83. Ashizawa T, Okada R, Suzuki Y, Takagi M, Yamazaki T, Sumi T, *et al.* Clinical significance of interleukin-6 (IL-6) in the spread of gastric cancer: Role of IL-6 as a prognostic factor. *Gastric Cancer* 2005;8:124-31.
84. Rojano-Rodríguez ME, Valenzuela-Salazar C, Cárdenas-Lailson LE, Romero Loera LS, Torres-Olalde M, Moreno-Portillo M. C-Reactive protein level in morbidly obese patients before and after bariatric surgery. *Revista de Gastroenterología de México (English Edition)* 2014;79:90-5.
85. Shimura T, Kitagawa M, Yamada T, Ebi M, Mizoshita T, Tanida S, *et al.* C-reactive protein is a potential prognostic factor for metastatic gastric cancer. *Anticancer Res* 2012;32:491-6.
86. Chen Z, Sun Y, Wang J, Shen X, Chen L, Zhuang Z. Prognostic and predictive value of serum C-reactive protein levels in patients with metastatic or locally recurrent gastric cancer. *Int J Biol Markers* 2016;31:e294-9.
87. Lee AV, Jackson JG, Gooch JL, Hilsenbeck SG, Coronado-Heinsohn E, Osborne CK, *et al.* Enhancement of insulin-like growth factor signaling in human breast cancer: Estrogen regulation of insulin receptor substrate-1 expression *in vitro* and *in vivo*. *Mol Endocrinol* 1999;13:787-96.
88. Clegg DJ, Brown LM, Woods SC, Benoit SC. Gonadal hormones determine sensitivity to central leptin and insulin. *Diabetes* 2006;55:978-87.
89. Edelstein ZR, Farrow DC, Bronner MP, Rosen SN, Vaughan TL. Central adiposity and risk of Barrett's esophagus. *Gastroenterology* 2007;133:403-11.
90. Lu Y, Lu F, Zeng S, Sun S, Lu L, Liu L. Genetics and gastric cancer susceptibility. *Int J Clin Exp Med* 2015;8:8377-83.
91. Motoshima H, Wu X, Sinha MK, Hardy VE, Rosato EL, Barbot DJ, *et al.* Differential regulation of adiponectin secretion from cultured human omental and subcutaneous adipocytes: Effects of insulin and rosiglitazone. *J Clin Endocrinol Metab* 2002;87:5662-7.
92. Hemminki K, Li X, Sundquist J, Sundquist K. Obesity and familial obesity and risk of cancer. *Eur J Cancer Prev* 2011;20:438-43.
93. Gulati P, Cheung MK, Antrobus R, Church CD, Harding HP, Tung YC, *et al.* Role for the obesity-related FTO gene in the cellular sensing of amino acids. *Proc Natl Acad Sci U S A* 2013;110:2557-62.
94. Vongsuvan R, George J, Qiao L, Van Der Poorten D. Visceral adiposity in gastrointestinal and hepatic carcinogenesis. *Cancer Lett* 2013;330:1-10.