

The Effects of 5-Aza-2'-Deoxycytidine and Valproic Acid on Apoptosis Induction and Cell Growth Inhibition in Colon Cancer HT 29 Cell Line

Abstract

Background: Epigenetic changes, including DNA methylation and histone modification, alter gene expression without the nucleotide template alterations and are associated with all stages of tumor formation and progression. Previously, we investigated the effects of DNA demethylating agents and histone deacetylase inhibitors on hepatocellular carcinoma and colon cancers. The current study aimed to investigate the effects of 5-aza-2'-deoxycytidine (5-AZA-CdR, decitabine) and valproic acid (VPA), individually as well as combined on apoptosis induction and cell growth inhibition in colon cancer HT 29 cell line. **Methods:** The effect of the compounds on the cell viability was measured by MTT assay. To determine cell apoptosis, the cells were treated with 5-aza-CdR and VPA. Propidium iodide was used for staining and the cells were analyzed using flow cytometry. **Results:** Both agents decreased cell viability in a time and dose-dependent manner significantly ($P < 0.002$). The results of flow cytometry demonstrated that 5-aza-CdR and VPA induced apoptosis significantly as opposed to control groups. Maximal percentage of apoptotic cells was obtained after 48 h with combined treatment. **Conclusions:** Our findings suggest that 5-aza-CdR and VPA can significantly inhibit cell growth and induce apoptosis in colon cancer HT 29 cell line.

Keywords: Apoptosis, colonic neoplasms, decitabine, valproic acid

Introduction

Colon cancer is the fourth leading cause of cancer deaths in the world.^[1] Cellular pathway disruptions that control cell proliferation and cell death play an important role in tumorigenesis, initiation, and progression of cancer. Epigenetic changes including DNA methylation and histone modification, alter gene expression without the nucleotide template alterations and are associated with all stages of tumor formation and progression.^[2] One of the best-known epigenetic changes that inactivate tumor suppressor genes is promoter CpG island hypermethylation mediated by DNA methyltransferases (DNMTs). Of all epigenetic changes, DNA hypermethylation, which inactivates transcription of tumor suppressor genes (TSGs) leading to gene silencing, has been reported in many cancers including colorectal, prostate, pancreatic, bladder, ovarian, and breast cancers.^[3] In mammalian cells, the DNMTs are divided into two major families, according to the structure and functions, including

maintenance methyltransferase (DNMT1) and de novo methyltransferases (DNMT3a, DNMT3b, and DNMT3L). Furthermore, DNMT2 displays a weak DNA methyltransferase activity. The high expression level of DNMT1 seen in the S-phase of the cell cycle makes it a target for DNMT inhibitors (DNMTIs) in cancer treatment.^[4] Other chromatin modifications associated with silenced TSGs are histone modifications such as acetylation and deacetylation. The histone acetylation levels are the result of the balance between histone deacetylase (HDAC) and histone acetyl-transferase (HAT) activities. HDACs remove the acetyl groups resulting in chromatin compaction. Mammalian HDACs can be subdivided into different groups according to their homology with yeast HDACs including class I (HDACs 1, 2, 3, and 8) and class II which are further subdivided into two subclasses: subclass IIa (HDAC 4, 7, and 9) and IIb (HDAC 6 and 10).^[5]

In vitro studies have revealed that DNA demethylating agents are involved in the reactivation of the silenced genes such

Masumeh Sanaei,
Fraidoon Kavooosi,
Hamed Sahraeian¹

Research Center for
Non-communicable Diseases,
Jahrom University of Medical
Sciences, Jahrom, Fars
Province, Iran, ¹Student of
Research Committee, Jahrom
University of Medical Sciences,
Jahrom, Fars Province, Iran

Address for correspondence:

Dr. Fraidoon Kavooosi,
Jahrom University of
Medical Sciences, Jahrom,
Fars Province, Iran.
E-mail: kavooosifraidoon@gmail.
com

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.ir

DOI:
10.4103/ijpvm.IJPVM_410_19

Quick Response Code:



How to cite this article: Sanaei M, Kavooosi F, Sahraeian H. The effects of 5-Aza-2'-deoxycytidine and valproic acid on apoptosis induction and cell growth inhibition in colon cancer HT 29 cell line. *Int J Prev Med* 2021;12:33.

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

as p16 in bladder, colon and pancreatic cells, p15 in AML cells and p57 in myeloid leukemia cells, and also re-expression of the cell cycle and apoptosis modulator Ras association domain family 1 isoform A (RASSF1A) in ovarian cancer cells.^[6] Natural DNMT inhibitor compounds such as curcumin, genistein, EGCG, resveratrol, equol, and parthenolide act through DNMTs inhibition.^[7] DNMT inhibitor 5-Aza-2'-deoxycytidine (5-aza-CdR) has strong potential to inhibit DNMT resulting in a progressive loss of methylation and reactivation of gene expression.^[8]

It is a prodrug that requires activation through phosphorylation by deoxycytidine kinase, which catalyzes the phosphorylation of deoxycytidine to form deoxycytidine monophosphate. The nucleotide analog, 5-aza-CdR, is incorporated into DNA and produces an irreversible inactivation of DNA methyltransferase. This compound is an S-phase specific agent that reactivates silent tumor-suppressor genes (TSGs) by DNA demethylation resulting in cell differentiation, cell growth inhibition, and cell apoptosis.^[9] HDACs can be divided into four groups structurally including hydroxamic acids, small molecular weight carboxylates, cyclic peptides, and benzamides. There are more than 15 HDACs some of which include SAHA (vorinostat), phenyl-butyrate, valproic acid (VPA), trapoxin A, and trichostatin A.^[10]

In cells cultured with HDACs, these compounds can upregulate the expression of both death receptors and their ligands in transformed cells. These agents upregulated proapoptotic proteins of the Bcl-2 family such as Bax, Bak, Bim, Bmf, and Bik and decrease Bcl-X_L, Bcl-w, and Mcl-1. The low concentration of HDACs induces G₁ arrest while high concentrations induce both G₁ and G₂/M arrests associated with induction of p21, which inhibits CDKs regulating G₁/S transition (CDK2), and G₁ progression (CDK4/6).^[11]

VPA, an 8-carbon branched-chained fatty acid has been reported as a well-known anticancer agent belonging to the small molecular weight carboxylates. This compound causes hyperacetylation of the N-terminal tails of histones H3 and H4 *in vivo* and *in vitro* and inhibits HDAC activity, it binds to the catalytic center thereby blocking substrate access.^[12] Previously, we investigated the effects of DNA demethylating agents and histone deacetylase inhibitors on hepatocellular carcinoma and colon cancers which encouraged us to design this study.^[13-20] The current study aimed to investigate the effects of 5-aza-CdR and VPA on apoptosis induction and cell growth inhibition in colon cancer HT 29 cell line.

Methods

Colon cancer HT 29 cells were purchased from the National Cell Bank of Iran, Pasteur Institute and were maintained in Dulbecco's modified Eagle's medium (DMEM). Culture media contains 100 mL/L fetal bovine serum (FBS)

and antibiotics (100 U/mL penicillin and 100 U/mL streptomycin) at 37°C in a humidified atmosphere of 5% carbon dioxide (CO₂). In addition, 5-aza-CdR and VPA were provided from Sigma (St. Louis, MO, USA) and dissolved in culture medium and phosphate-buffered saline (PBS), respectively at a final concentration of 100 mM to prepare a suitable stock solution. All of the other test concentrations were prepared by diluting the stock solution. DMEM, FBS, and 3-[4,5-dimethyl-2-thiazolyl]-2,5-diphenyl-2H-tetrazolium bromide (MTT) were obtained from Sigma. Dimethyl sulfoxide (DMSO) was purchased from Merck Co. (Darmstadt, Germany).

Cell viability assay

The effect of 5-aza-CdR and VPA on the cell viability was measured by MTT assay. The cells were seeded into the 96-well plates at 5×10^5 cells/well and allowed to adhere overnight. After 24 h, the culture medium was replaced with medium containing different concentrations of VPA (0.5, 1, 2.5, 5, and 10 μ M) and 5-aza-CdR (0.5, 1, 2.5, 5, and 10 μ M), except control groups which were treated with a drug-free medium for different periods. After treatment times (24 and 48 h), the viable cells were determined by an MTT assay, as described elsewhere.^[11]

Apoptosis assay

The colon cancer HT 29 cell, 5×10^5 cells, were treated with 5-aza-CdR at a concentration of 2 μ M and VPA at a concentration of 5 μ M and the cells were harvested by trypsin solution at the end of the treatment (after 24 and 48 h). Besides, one group was treated with 5-aza-CdR at a concentration of 2 μ M for 24 h followed by VPA at a concentration of 5 μ M for another 24 h. After treatment times, the cells were washed with cold phosphate-buffered saline (PBS) and resuspended in binding buffer (1x). Finally, Annexin-V-(FITC) and propidium iodide (PI, Becton-Dickinson, San Diego, CA) were used for staining according to the manufacturer's instructions and the cells were analyzed using flow cytometry.

Statistical analysis

The data were obtained from three tests and are shown as means \pm standard deviations. Statistical comparisons between groups were performed with ANOVA (one-way ANOVA) and Turkey test. A significant difference was considered as $P < 0.05$.

Results

Result of cell viability assay

The colon cancer HT 29 cell were cultured in the 96-well plates and treated with VPA (0.5, 1, 2.5, 5, and 10 μ M) and 5-aa-CdR (0.5, 1, 2.5, 5, and 10 μ M) for 24 and 48 h and evaluated by MTT assay to determine cell viability. The results are shown in Figure 1, comparing the percentage of cell viability in treated cells with untreated control

groups. As shown in Figure 1, both compounds decreased cell viability in a time and dose-dependent manner significantly ($P < 0.002$). The IC₅₀ values of 5-aza-CdR and VPA were 2 μ M and 5 μ M, respectively.

Result of apoptosis assay

To determine the apoptotic effect of the compounds, the propidium iodide (PI) and Annexin-V/FITC staining were done. The results demonstrated that 5-aza-CdR at a concentration of 2 μ M and VPA at a concentration of 5 μ M induced apoptosis significantly as opposed to control groups. A similar result was observed in the group treated with 5-aza-CdR at a concentration of 2 μ M for 24 h followed by VPA at a concentration of 5 μ M for another 24 h. The percentage of apoptotic cells in 5-aza-CdR-treated groups was 47 and 82% and in VPA-treated groups were 85 and 94%, respectively ($P < 0.001$). The percentage of apoptotic cells in the group which treated with 5-aza-CdR at a concentration of 2 μ M for 24 h and followed by VPA at a concentration of 5 μ M for another 24 h were 96% ($P < 0.001$). Maximal percentage of apoptotic cells was obtained after 48 h with combined treatment [Figures 2 and 3].

Discussion

Colon cancer is a leading cause of cancer deaths worldwide. One of the basic causes of the colon cancer initiation and progression is the accumulation of a variety of epigenetic changes such as aberrant DNA methylation and histone deacetylation in colon epithelial cells. Epigenetic assessment of colon cancer has indicated aberrantly methylated and histone deacetylated genes that alter miRNA expression.^[21] DNA hypermethylation-mediated epigenetic silencing of TSGs has emphasized the development and application of anticancer drugs which work by inhibition of DNA methylation and restore silenced genes by reprogramming the genes involved in cancer initiation and progression. DNMTs modulate the cellular epigenome by reversing the hypermethylation, resulting in renewed transcription of silenced TSGs.^[22]




Figure 1: Effect of VPA (0.5, 1, 2.5, 5, and 10 μ M) and 5-aza-CdR (0.5, 1, 2.5, 5, and 10 μ M) on colon cancer HT 29 cell viability determined by the MTT assay. Data are presented as mean \pm SD from at least triplicate wells and three independent experiments. Both compounds with all concentrations indicated a significant inhibitory effect. The first column of each group belongs to the control group

HDACIs are natural or synthetic agents that have broad functions in cancer treatment. These compounds can change the balance between HATs and HDACs, resulting in the accumulation of acetylated histones/nonhistone proteins, which reactivates silenced TSGs.^[23] Previously, we reported the effect of DNA demethylating agents such as genistein and histone deacetylase inhibitors such as VPA on cancers.^[13-17] The result of the current study indicated that 5-aza-CdR and VPA (individually and combined) can inhibit cell growth and induce significant apoptosis in colon cancer cells. One of the major mechanisms of DNA demethylating agents and histone deacetylase inhibitors is silenced TSGs reactivation as we reported estrogen receptor alpha (ER α) reactivation and DNMTs suppression previously. It has been reported that 5-aza-CR down-regulates DNMT1, DNMT3a, and DNMT3b gene expression, arrests the cell cycle at the G2/M phase and induces apoptosis by a joined activation of p53 protein and its downstream effectors p21WAF1 and GADD45 in the colon cancer cell line HCT-116.^[24] Other studies have demonstrated that the 5-aza-CdR treatment affected the cell cycle and caused accumulation of cells in the G0/G1 phase by up-regulating the silenced tumor suppressor Ras association domain family 1A (RASSF1A) gene in Caco-2 cells.^[25] Furthermore, it has been shown that treatment with 5-aza-CdR up-regulates endothelial PAS domain-containing protein 1 (EPAS1) expression in




Figure 2: Apoptotic effects of VPA and 5-aza-CdR. Both compounds induced significant apoptosis in colon cancer HT 29 cells. The apoptotic cell percentage in the group treated with 5-aza-CdR (24 h) and then VPA (24 h) were more significant than that of each drug alone. ($P < 0.001$ compared with control group)




Figure 3: Apoptotic effects of VPA and 5-aza-CdR on colon cancer HT 29 cell. Combined drug treatment-induced apoptosis more significant than each drug alone ($P < 0.001$). Asterisks (*) indicate significant differences between treated cells and the control group

colorectal cancer cells which are associated with DNA demethylation of the EPAS1 regulatory region.^[26] In agreement with our report, it has been reported that VPA inhibits cell growth and induces apoptosis in the HT-29 and LoVo colon cancer cell lines. It modulates the expression of PPAR γ , a possible heterodimer of RXRs. VPA exerts its effect by modulation of the pro- and antiapoptotic factors, BAX and BCL2, respectively, it enhances the expression of BAX and down-regulates BCL2 expression in the HT-29 and LoVo colon cancer cell lines.^[27] VPA blocks cell growth, possibly through the synthesis of p21WAF1/CIP1 and p27KIP1. It can exert by two main pathways resulting in programmed cell death including death receptor (extrinsic) pathway and the mitochondrial (intrinsic) pathway. Activation of both caspase-8 and caspase-9 has been demonstrated in VPA-treated colorectal cells.^[28] Based on our result, combined treatment indicated a more significant apoptotic effect than that of a single agent alone. In constant with our finding, it has been reported that the combination of 5-aza-CdR and VPA has a synergistic effect in terms of cell growth inhibition, apoptosis induction, and reactivation of p57KIP2 and p21CIP1 in leukemic cell lines HL-60 and MOLTA.^[29] Other investigators have shown that DNA methyltransferase/histone deacetylase inhibitor combinations reverse the aberrant epigenetic gene silencing which leads to apoptotic induction in myeloid neoplasms.^[30] Furthermore, histone deacetylase inhibitor trichostatin A can significantly upregulate DNA methylase inhibitor 5-aza-CdR -induced MAGE-A1, -A2, -A3, and -A12 genes expression in breast cancer (MDA-MB-231, GI101, and MTSV1.7) and colon cancer (WiDr) cell lines.^[31] In all, our finding indicated that 5-aza-CdR and VPA could induce apoptosis in colon cancer HT 29 cell line, if used separately or in combination.

Conclusions

In conclusion, our finding shows that 5-aza-CdR and VPA can inhibit *in vitro* colon cancer HT 29 cells and induce

their apoptosis. Our results provide experimental support and a new therapeutic strategy into novel therapies for colon cancer. Meanwhile, the current study is limited to *in vitro* experiments, hence further evaluations will be required for a better understanding of the effects of these compounds on colon cancer.

Acknowledgments

This article was supported by the adjutancy of research of Jahrom University of Medical Sciences, Iran. This article has been extracted from Ms. Hamed Sahraeian's thesis.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Received: 29 Oct 19

Accepted: 25 Dec 19

Published: 29 Mar 21

References

- Ng J, Yu J. Promoter hypermethylation of tumour suppressor genes as potential biomarkers in colorectal cancer. *Int J Mol Sci* 2015;16:2472-96.
- Grønbaek K, Treppendahl M, Asmar F, Guldborg P. Epigenetic changes in cancer as potential targets for prophylaxis and maintenance therapy. *Basic Clin Pharmacol Toxicol* 2008;103:389-96.
- Heichman KA, Warren JD. DNA methylation biomarkers and their utility for solid cancer diagnostics. *Clin Chem Lab Med* 2012;50:1707-21.
- Delpu Y, Cordelier P, Cho W, Torrisani J. DNA methylation and cancer diagnosis. *Int J Mol Sci* 2013;14:15029-58.
- Ropero S, Esteller M. The role of histone deacetylases (HDACs) in human cancer. *Mol Oncol* 2007;1:19-25.
- Gnyszka A, Jastrzebski Z, Flis S. DNA methyltransferase inhibitors and their emerging role in epigenetic therapy of cancer. *Anticancer Res* 2013;33:2989-96.
- Singh V, Sharma P, Capalash N. DNA methyltransferase-1 inhibitors as epigenetic therapy for cancer. *Curr Cancer Drug Targets* 2013;13:379-99.
- Robertson KD, A. Jones P. DNA methylation: Past, present and future directions. *Carcinogenesis* 2000;21:461-7.
- Momparler RL. Epigenetic therapy of cancer with 5-aza-2'-deoxycytidine (decitabine). *Seminars in Oncology*. 32. Elsevier; 2005. p. 443-51.
- Federico M, Bagella L. Histone deacetylase inhibitors in the treatment of hematological malignancies and solid tumors. *Biomed Res Int* 2010;2011:1-12.
- Xu W, Parmigiani R, Marks P. Histone deacetylase inhibitors: Molecular mechanisms of action. *Oncogene* 2007;26:5541.
- Chavez-Blanco A, Perez-Plasencia C, Perez-Cardenas E, Carrasco-Legleu C, Rangel-Lopez E, Segura-Pacheco B, *et al.* Antineoplastic effects of the DNA methylation inhibitor hydralazine and the histone deacetylase inhibitor valproic acid in cancer cell lines. *Cancer Cell Int* 2006;6:2.
- Kavoosi F, Sanaei M. Comparative analysis of the effects of valproic acid and tamoxifen on proliferation, and apoptosis of human hepatocellular carcinoma WCH 17 CellLin. *Iran J Ped Hematol Oncol* 2018;8:12-20.

14. Sanaei M, Kavooosi F, Mansoori O. Effect of valproic acid in comparison with vorinostat on cell growth inhibition and apoptosis induction in the human colon cancer SW48 cells *in vitro*. *Exp Oncol* 2018;40:95-100.
15. Sanaei M, Kavooosi F, Roustazadeh A, Shahsavani H. *In vitro* effect of the histone deacetylase inhibitor valproic acid on viability and apoptosis of the PLC/PRF5 human hepatocellular carcinoma cell line. *Asian Pac J Cancer Prev* 2018;19:2507-10.
16. Sanaei M, Kavooosi F, Salehi H. Genistein and trichostatin A induction of estrogen receptor alpha gene expression, apoptosis and cell growth inhibition in hepatocellular carcinoma HepG2 cells. *Asian Pac J Cancer Prev* 2017;18:3445.
17. Sanaei M, Kavooosi F, Atashpour S, Haghighat S. Effects of genistein and synergistic action in combination with tamoxifen on the HepG2 human hepatocellular carcinoma cell line. *Asian Pac J Cancer Prev* 2017;18:2381.
18. Sanaei M, Kavooosi F, Pourahmadi M, Moosavi SN. Effect of genistein and 17- β estradiol on the viability and apoptosis of human hepatocellular carcinoma HepG2 cell line. *Adv Biomed Res* 2017;6:163.
19. Sanaei M, Kavooosi F, Roustazadeh A, Golestan F. Effect of Genistein in comparison with trichostatin A on reactivation of DNMTs genes in hepatocellular carcinoma. *J Clin Transl Hepatol* 2018;6:1-6.
20. Sanaei M, Kavooosi F, Valiani A, Ghobadifar MA. Effect of genistein on apoptosis and proliferation of hepatocellular Carcinoma Hepa1-6 Cell Line. *Int J Prev Med* 2018;9:12.
21. Okugawa Y, Grady WM, Goel A. Epigenetic alterations in colorectal cancer: Emerging biomarkers. *Gastroenterology* 2015;149:1204-25. e12.
22. Agrawal K, Das V, Vyas P, Hajdúch M. Nucleosidic DNA demethylating epigenetic drugs—A comprehensive review from discovery to clinic. *Pharmacol Ther* 2018;188:45-79.
23. Chen HP, Zhao YT, Zhao TC. Histone deacetylases and mechanisms of regulation of gene expression. *Crit Rev Oncog* 2015;20:35-47.
24. Schneider-Stock R, Diab-Assef M, Rohrbeck A, Foltzer-Jourdainne C, Boltze C, Hartig R, *et al.* RETRACTION: 5-aza-cytidine is a potent inhibitor of DNA methyltransferase 3a and induces apoptosis in HCT-116 colon cancer cells via Gadd45-and p53-dependent mechanisms. *J Pharmacol Exp Ther* 2005;312:525-36.
25. Li X, Qin B, Liu B. Delineating the effect of demethylating agent 5-aza-2'-deoxycytidine on human Caco-2 colonic carcinoma cells. *Oncol Lett* 2016;12:139-43.
26. Rawluszko-Wieczorek AA, Horbacka K, Krokowicz P, Misztal M, Jagodziński PP. Prognostic potential of DNA methylation and transcript levels of HIF1A and EPAS1 in colorectal cancer. *Mol Cancer Res* 2014;12:1112-27.
27. Papi A, Ferreri A, Guerra F, Orlandi M. Anti-invasive effects and proapoptotic activity induction by the rexinoid IIF and valproic acid in combination on colon cancer cell lines. *Anticancer Res* 2012;32:2855-62.
28. Mologni L, Cleris L, Magistroni V, Piazza R, Boschelli F, Formelli F, *et al.* Valproic acid enhances bosutinib cytotoxicity in colon cancer cells. *Int J Cancer* 2009;124:1990-6.
29. Yang H, Hoshino K, Sanchez-Gonzalez B, Kantarjian H, Garcia-Manero G. Antileukemia activity of the combination of 5-aza-CdR with valproic acid. *Leukemia Res* 2005;29:739-48.
30. Gore SD, Baylin S, Sugar E, Carraway H, Miller CB, Carducci M, *et al.* Combined DNA methyltransferase and histone deacetylase inhibition in the treatment of myeloid neoplasms. *Cancer Res* 2006;66:6361-9.
31. Wischniewski F, Pantel K, Schwarzenbach H. Promoter demethylation and histone acetylation mediate gene expression of MAGE-A1,-A2,-A3, and-A12 in human cancer cells. *Mol Cancer Res* 2006;4:339-49.