

Systematic Review and Meta-Analysis on Quality of Life in Diabetic Patients in Iran

Abstract

Background: Diabetes is the fifth leading cause of death in the world, which reduces the patients' quality of life (QOL) and is considered as an important subject especially in medicine and medical community. The present study aimed at investigating the QOL of diabetic patients in Iran through meta-analysis. **Methods:** The search was conducted using relevant keywords in national and international databases including Iranmedex, SID, Magiran, IranDoc, Medlib, Science Direct, PubMed, Scopus, Cochrane, Embase, Web of Science. Questionnaires WHOQOL, SF-36, SF-20, DQOL, QOL, PedsQL, ADDQOL, D-39, DQOL-BCI, SWED-QUAL, IRDQOL, PHG-2, EQ-5D, and IDQOL-BCI were used to assess the QOL. Heterogeneity of studies was assessed using I² index. Data were analyzed using STATA version 11. **Results:** In 96 studies of 17,994 people, the mean score of QOL in diabetic patients was based on the questionnaires WHOQOL [66.55 (95% CI: 45.83, 87.26)], D-39 [129.43 (95%CI: 88.77, 170.10)], SF-36 [65.64 (95% CI: 59.82, 71.46)], SF-20 [46.50 (95% CI: 37.19, 55.81)], DQOL [61.19 (95% CI: 35.73, 86.66)], QOL [117.91 (95% CI: -62.97, 298.79)], PedsQL [34.36 (95% CI: -31.49, 100.22)], ADDQOL [41.76 (95% CI: 12.01-71.50)], SWED-QUAL [59.19 (95% CI: 21.15, 97.23)], IRDQOL [105.92 (95% CI: 102.73, 109.10)], PHG-2 [61.00 (95%CI: 59.63, 62.37)], EQ-5D [0.62 (95% CI: 0.61, 0.64)], DQOL-BCI [3.40 (95% CI: 3.31, 3.49)], and IDQOL-BCI [22.63 (95% CI: -2.38, 47.64)]. **Conclusions:** The QOL of diabetic patients was evaluated according to different types of questionnaires and the QOL of diabetic patients was found to be lower than normal population.

Keywords: Diabetes, meta-analysis, quality of life, systematic review

Introduction

World Health Organization defines quality of life (QOL) as an individuals' understanding of living condition in terms of culture and the prevailing community values following their goals, expectations, standards, and interests. Hence, QOL is closely related to physical, psychological, and mental condition, personal beliefs, level of self-reliance, mass communication, and environment.^[1-6] One reason for the multidimensional complexity of QOL is that it includes different aspects of an individual's life. Another reason is that each individual has his/her own unique characteristics and his/her perception of a good or poor QOL is unique to that person.^[1-9] The subject of QOL is important since it may lead to frustration, lack of motivation for any attempt and reduction of social, economic, cultural,

and health activities. QOL influences the socioeconomic development of a country in deeper dimensions. Modifying the QOL is considered as a part of disease control program.^[7-9]

Diabetes is known as a "silent epidemic" and is considered a major public health problem in the United States and other parts of the world, including Iran. It is the most prevalent metabolic disease with an increasing incidence, which shortens life expectancy by one third^[10-12] and affects various aspects of a patient's life, including psychological, physical, social, and economic condition, family life and sexual function.^[13-16] Type 1 and type 2 diabetes are two major forms of this disease and include about 10--90% of the diabetes population, respectively.^[17]

1. According to the latest available data, about 171 million people suffer from diabetes worldwide. Asia is one of the regions with a high prevalence of diabetes.^[18] Two percent of the Iranian population are suffering from

Moloud Fakhri,
Mohsen Abdan¹,
Melina Ramezanpour²,
Ali Hasanpour
Dehkordi^{3,*},
Diana Sarikhani^{1,*}

Traditional and Complementary
Medicine Research Center,
Addiction Institute, Mazandaran
Sari, University of Medical
Sciences, ²Medical Student at
school of medicine, Mazandaran
University of Medical Sciences,
Sari, ¹Research Center for
Environmental Determinants
of Health (RCEDH), School
of Public Health, Kermanshah
University of Medical
Sciences, Kermanshah, ³Social
Determinants of Health
Research Center, School of
Allied Medical Sciences,
Shahrekord University of
Medical Sciences, Shahrekord,
Iran

Address for correspondence:
Dr. Diana Sarikhani,
Research Center for
Environmental Determinants
of Health (RCEDH), School
of Public Health, Kermanshah
University of Medical
Sciences, Kermanshah, Iran.
E-mail: diana_sarokhani@yahoo.com
Dr. Ali Hasanpour Dehkordi
Social Determinants of Health
Research Center, School of
Allied Medical Sciences,
Shahrekord University of
Medical Sciences, Shahrekord,
Iran

1 Quality of life

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-Non Commercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

How to cite this article: Fakhri M, Abdan M, Ramezanpour M, Dehkordi AH, Sarikhani D. Systematic review and meta-analysis on quality of life in diabetic patients in Iran. *Int J Prev Med* 2021;12:41.

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.net

DOI:
10.4103/ijpvm.IJPVM_327_19

Quick Response Code:



the disease.^[19] Due to the large proportion of diabetic patients in Iran and the direct impact of diabetes on the QOL of patients with diabetes, the present study aims to evaluate the QOL in diabetic patients in Iran. Considering that a meta-analysis study of the same title was published in 2016^[20] and evaluated only two questionnaires (SF-20 and SF-36). Also the previous meta-analysis included only the results of 10 studies. The present meta-analysis was performed with the aim of updating the previous study and without considering the time limit, limiting the type of questionnaire and covering all studies published in this field. In the present meta-analysis, the QOL of diabetic patients was evaluated in the form of levels: Good, Moderate, and Poor. This issue was not presented in previous meta-analysis.

Methods

Search strategy

This is a systematic review and meta-analysis aimed at investigating the QOL of diabetic patients in Iran. In order to achieve the related documentation in Persian and English, two researchers independently searched both national and international databases, including Iranmedex, SID, Magiran, Iran-Doc, Med-Lib, Science-Direct, PubMed, Scopus, Cochrane, Embase, Web of Science, and Medline using related Persian keywords and their English equivalents: "Iran," "meta-analysis," "diabetes," and "quality of life." The keywords were searched using AND/OR operators. The search was performed without time limit until 22.04.2020. However, the articles in question were published between 2003 and 2020. The previous meta-analysis article published in this field belonged to 2016^[20] and only examined the SF-20 and SF-36 questionnaires, while the current meta-analysis did not impose any restrictions on the type of questionnaires used in the reviewed articles. For this reason, various questionnaires such as: WHOQOL, SF-36, SF-20, DQOL, PedsQL, ADDQOL, D-39, DQOL-BCI, SWED-QUAL, and IRDQOL were evaluated. In cases of lack of access to the article's full text, the researchers asked the corresponding author for the full-text articles via email. To complete the search, Google Scholar was also searched.

Inclusion and exclusion criteria

Inclusion criteria included mentioning the QOL of diabetic patients in Iran in Persian and English. Exclusion criteria included non-random sampling, inadequate information in the article's text, and population other than diabetic patients.

Study selection

In the first phase of the search, 501 articles related to QOL of diabetic patients were found. After reviewing the titles, 289 duplicate and overlapping articles were

excluded. Abstracts of all remaining articles were reviewed and 59 irrelevant articles were excluded. The full text of the remaining articles was reviewed then 57 studies were excluded due to having the exclusion criteria. Finally, 96 articles entered the qualitative evaluation stage [Chart 1].

Qualitative evaluation of studies

To check the quality of studies, the STROBE checklist (strengthening the reporting of observational studies in epidemiology)^[21] was applied. This checklist includes 22 items that cover different parts of a report (sampling, measuring variables, objectives of the study, and statistical analysis). Each item was given one point and higher points were given to other items that we considered more important. In this phase, four unqualified articles were excluded and finally 96 articles entered the meta-analysis stage.

Data extraction

To reduce bias in reporting and error in data collection, two researchers independently extracted data from articles and entered the data into a checklist, which included the following items: The first author's name, title of study, sample size, year of publication, city of study, diabetes type, questionnaire title, the subjects' average age, mean and standard deviation of the QOL of diabetic patients, mean and standard deviation of quality of life dimensions, etc.

Statistical analysis

Considering that the QOL in diabetic patients score and its subgroups score were quantitative, the mean and standard deviation of these indices were extracted in each study and the variance of the mean was calculated using normal distribution. Considering the heterogeneity of the studies, a random effects model was used to combine the results of the studies. The I^2 index was used to investigate the heterogeneity of the studies. A random effects meta-analysis was used to give a pooled estimate of prevalence of QOL for each measure. Metaregression was used to check heterogeneity




Figure 1: The relationship between quality of life score in diabetic patients and the year of publication




Chart 1: Flowchart of the entrance of studies into the systematic review and meta-analysis

among the studies and to find any association between the year of publication and the sample size with QOL in diabetic patients. Subgroup analysis was done according to sex, components, and questionnaire. All statistical analyses were performed using STATA ver 14. The significance level of the tests was considered to be $P < 0.05$.

Results

In 96 reviewed studies with a sample of 17,994, the mean QOL score in diabetic patients was based on WHO Quality of Life-BREF (WHOQOL-BREF) [66.55 (95% CI: 45.83, 87.26)], D-39 [129.43 (95% CI: 88.77, 170.10)], *Short Form-36* (SF-36) [65.64 (95% CI: 59.82, 71.46)], *Short Form-20* (SF-20) [46.50 (95% CI: 37.19, 55.81)], Diabetes Quality of Life (DQOL) [61.19 (95% CI: 35.73, 86.66)], Quality of Life [QOL] (117.91 (95% CI: -62.97, 298.79)], PedsQL [34.36 (95% CI: -31.49, 100.22)], Audit of Diabetes Dependent Quality of Life (ADDQOL) [41.76 (95% CI: 12.01-71.50)], SWED-QUAL [59.19 (95% CI: 21.15, 97.23)], IRDQOL [105.92 (95% CI: 102.73, 109.10)], PHG-2 [61.00 (95% CI: 59.63, 62.37)], EQ-5D [0.62 (95% CI: 0.61, 0.64)], DQOL-BCI [3.40 (95% CI: 3.31, 3.49)], and Iranian version of the Diabetes Quality of Life Brief Clinical Inventory (IDQOL-BCI) [22.63 (95% CI: -2.38, 47.64)]. Considering the heterogeneity between the studies, the confidence interval for each study based on random-effects model is shown in Table 1.

According to the results, the mean QOL score in diabetic patients is presented in Table 2. In the WHOQOL-BREF2 questionnaire, the highest and lowest scores of QOL score in diabetic patients were related to **Social Activity** (48.36) and the **Mental** (36.29), respectively. In the SF-36

questionnaire, the highest and lowest quality of life scores of diabetic patients were related to **Limitation of Activity** (52.72) and **Peripheral** (24.10), respectively. The mental dimension (20.75) and the **Peripheral** (9.60) had the highest and lowest QOL scores of diabetic patients in the SF-203 questionnaire. In the DQOL4 questionnaire, the highest and lowest QOL scores of diabetic patients were related to **General Health** dimension (41.25) and **Social Activity** (13.46), respectively. In the QOL5 questionnaire, the **Peripheral** dimension (20.23) and the **Social Activity** dimension (5.18) had the highest and lowest QOL scores of diabetic patients, respectively. In the PedsQL6 questionnaire, the highest and lowest QOL scores of diabetic patients were related to **Emotion** dimension (59.84) and **Peripheral** dimension (33.15), respectively. In the SWED-QUAL questionnaire, the highest and lowest QOL scores of diabetic patients were related to **Physical** dimension (21.84) and **Physical Pain** dimension (8.07), respectively. In the IRDQOL questionnaire, the highest and lowest QOL scores of diabetic patients were related to **Social Activity** dimension (69.53) and **Physical** dimension (57.03), respectively. In the PHG-2 questionnaire, the highest and lowest QOL scores of diabetic patients were related to **Physical** dimension (16.43) and **Emotion** dimension (9.84), respectively.

In the Sf-36 questionnaire, 15% of diabetic patients had a good QOL and 46% had a poor QOL. In the Sf-20 questionnaire, 29% of diabetic patients had a good QOL and 36% had a low QOL. In the QOL questionnaire, 36% of diabetic patients had a desirable QOL and 45% had a poor QOL. In the WHOQOL questionnaire, 55% of diabetic patients had an acceptable QOL and 37% had a poor QOL. In the SWED-QUAL questionnaire, 62% of diabetic patients had an acceptable QOL and 38% had a poor QOL. In the IRDQOL questionnaire, 11% of diabetic patients had an acceptable QOL and 66% had a poor QOL [Table 3].

In order to perform additional analyzes, we plotted the meta-regression diagram. There was no significant statistical relationship in the study of meta-regression score of quality of life in diabetic patients based on the year of study ($P = 0.565$) [Figure 1]. This means that over time, the QOL of diabetic patients has not decreased. The relationship between QOL score in diabetic patients and the number of research samples was not statistically significant ($P = 0.106$) [Figure 2].

Discussion

In 96 reviewed studies with a sample of 17,994, the QOL score in diabetic patients was 66.55 in WHOQOL, 65.64 in SF-36, 46.50 in SF-20, 61.19 in DQOL, 117.91

- 3 Short Form
- 4 Diabetes Quality of Life
- 5 Quality of Life
- 6 Pediatric Quality of Life Inventory

2 WHO Quality of Life-BREF

Table 1: Specifications of reviewed articles on the status of the quality of life of diabetic patients in Iran

ID	Author	Year of publication	City of study	Type of diabet	Sample size	Age mean	Questionnaire	Mean score of QOL	SD of QOL
[22]	Aghamolaei T	2003	Hormozgan	type 2	80	32-72	WHOQOL-BREF		
[23]	Aghamolaei T	2005	Hormozgan	type 2	71	51.3	WHOQOL-BREF		
[24]	Sadeghie Ahari S	2008	Ardebil	type 2	110	52.5	SF-36		
[25]	Ahmadi A	2011	Chaharmahal& Bakhtiari	type 2	254	30-65	Developed by reserch team		
[26]	Alavi A	2010	Chaharmahal& Bakhtiari	type 1	22	15.33	PedsQL	0.78	0.48
[27]	Baghianimoghadam MH	2008	Yazd	type 2	120	25-75	SF-20	51.03	17.04
[28]	Bazzazian S	2010	Tehran	type 1	300	18-30	D-39	109.47	45.31
[29]	Borzou SR	2010	Hamedan	type 2	165		SF-36		
[30]	Safarabadi-Farahani T	2010	Tehran	type 1	70	14.94	DQOL for youth	56.28	12.2
[31]	Ghanbari A	2004	Guilan	type 2	90	>40	SWED-QUAL	18.37	12.5
[32]	Ghanbari A	2005	East-Azerbaijan	type 2	117	>35	SWED-QUAL	28	8.1
[33]	Haririan HR	2009	East-Azerbaijan	type 2	150	20-60	SWED-QUAL		
[34]	Heydari M	2007	Zanjan	type 1	47	11-20.	Developed by reserch team	106.65	45.75
[35]	Jafari P	2011	Fars	type 1	94	8-.18	PedsQL	67.98	14.03
[36]	Ghavami H	2005	west-Azerbaijan	type 2	74	40-65	Developed by reserch team	98	
[37]	Shahab-Jahanlou AR	2011	Hormozgan	type 2	256	27-72	WHOQOL-BREF26		
[38]	Shahab-Jahanlou AR	2011	Hormozgan	typ1& type 2	76	49.15	IRDQOL		
[1]	Darvishpour-Kakhaki A	2005	Tehran	typ1& type 2	131	47.3	SF-36		
[39]	Sedaghati-Kasbakhi M	2008	Mazandaran	type 2	70		SWED-QUAL	131.72	25.88
[40]	Kermansaravi F	2012	Sistan and Baluchestan	type 1	100	14.6	DQOL for youth	52.65	14.58
[41]	Khaledi S	2011	Kurdistan	type 2	198	>18	SF-36	70.82	18.97
[42]	Khamseh MA	2011	Tehran	type 1	150	22.14	Developed by reserch team	69.01	13.03
[43]	Peymani M	2007	Tehran	typ 1 and type 2	302	>18	Developed by reserch team		
[44]	Rakhshanderu S	2006	Tehran	type 2	40	40-65	DQOL	35.2	9.1
[45]	Rasouli D	2011	Tehran	patients with deiabetic foot ulcer	120	54.23	DFS		
[46]	Safavi M	2011	Ardebil	type 2	123	30-70	QOL	234.27	5.18
[47]	Sanjari M	2011	Kerman	typ 1 and type 2	132	52.98	SF-36	314.18	138.24
[48]	Shahrjerdi S	2009	Markazi	type 2	27	>35	SF-36	83.08	11.06
[49]	Sayadi N	2011	Khuzestan	type 2	31	58.35	SF-36	1775.81	955.4
[50]	Taghdisi MH	2011	Golestan	type 2	78	49	WHOQOL	80.39	11.35
[51]	Timareh M	2012	Kermanshah	typ 1 and type 2	350	>18	SF-36		
[52]	Vares Z	2010	Isfahan	typ 1 and type 2	310	>18	IRDQOL	105.8	44.1
[53]	Vazirinezhad R	2010	Kerman		101	50.8	SF-36		
[54]	Yekta Z	2011	West-Azerbaijan	type 2	250	60.73	SF-36	57.52	17.1
[38]	Shahab-Jahanlou AR	2011	Hormozgan	typ 1 and type 2	76	49.15	WHOQOL		
[55]	Mirfeizi M	2012	Karaj		180	53.47	IDQOL-BCI	9.89	2.51
[56]	Shahi M	2017	Semnan	type 2	60	57.82	QOL		
[57]	Najafi-Ghezeljeh T	2017	Tehran	type 2	65	54.3	IDQOL-BCI	35.41	7.8
[58]	Shamshirgaran SM	2016	Ardebil	type 2	300	54.13	WHOQOL	53.07	7.09
[59]	Hajian-Tilaki K	2016	Babol		750	67.85	SF-36		
[60]	Dadgostar H	2016	Tehran	type 2	74	49.65	SF-36		
[61]	Jafari N	2014	Isfahan	type 2	203	55.42	PHG-2	61	9.97
[62]	Abdoli S	2015	Malayer	type 2	40	35-85	WHOQOL-BREF		
[63]	Hadi N	2013	Shiraz	typ 1 and type 2	300	50.98	SF-36		

Contd...

Table 1: Contd...

ID	Author	Year of publication	City of study	Type of diabet	Sample size	Age mean	Questionnaire	Mean score of QOL	SD of QOL
[64]	Shavandi N	2010	Markazi	type 2	17	48.52	SF-36	74.58	11.34
[65]	Shayeghian Z	2013	Tehran	type 2	100	55.4	ADDQoL	26.63	12.01
[66]	Alipour A	2012	Yaza	type 2	80	46.2	ADDQoL	56.98	18.63
[67]	Afshar M	2014	Kashan	type 2	56	14.75	IRDQOL	106	15.95
[68]	Derakhshanpour F	2015	Gorgan	type 2	330	50.6	WHOQOL-BREF	54.79	13.7
[69]	Zaker MR	2016	Urmia		80		DQOL	46.04	4.3
[70]	Didarloo AR	2016	Khoy	type 2	352	43	WHOQOL-BREF	58.02	17.63
[71]	Gholami A	2013	Neishabour	type 2	1847	59.65	WHOQOL-BREF	12.18	2.3
[72]	Torabi M	2014	Hamedan	type 2	110	47.4	SF-36		
[73]	Izadi A	2014	Khoram Abad	type 2	80	30-70	SF-20		
[74]	Khodabakhsi-Kulaei A	2015	Tafresh	type 2	24	50.58	WHOQOL	68	11.08
[75]	Mohammad-Shahi A	2014	Ahvaz	type 2	110	53.69	SF-36		
[76]	Saeedpour J	2013	Tehran		60	40	SF-36	43.5	15.7
[4]	Masoudi-Alavi N	2004	Tehran	typ 1 and type 2	104	50.5	QOL	116.7	18.8
[77]	Ghasemipour M	2009	Khoram Abad		150	18-65	QOL	2.77	0.79
[78]	Eydi-Bayegi M	2014	Ahvaz	type 2	50	46.2	WHOQOL-26	73.91	14.85
[79]	Sadeghi T	2012	Rafsanjan		70	18-65	SF-36		
[80]	Zaree-Bahramabadi M	2012	Sanandaj	type 2	48	30-50	SF-36	53.3	10.76
[81]	Qashqaei S	2014	Shiraz	type 2	42	35-65	SF-36	56.37	18.25
[82]	Saadatjuo SAR	2012	Birjand	type 2	100	42.82	SF-36	57.29	26.09
[83]	Behrooz B	2016	Kermanshah	type 2	16	49.47	WHOQOL-26	137.92	12.9
[84]	Ebrahimi H	2014	Shahrood	type 2	156	48.11	DQOL	164.53	63.21
[85]	Mohammadshahi GHR	2016	Taybad	type 2	20	47.75	SF-36		
[86]	Shams S	2015	Urmia		80		SF-36		
[87]	Mohammadpour Y	2008	Tabriz	type 2	150		Self-made		
[88]	Ganjluo J	2015	Sabzevar	type 2	75	35-65	ADDQOL-19		
[89]	Bidi F	2012	Bojnord	type 2	40	52.17	SF-20	41.52	16.28
[90]	Derakhshanpour F	2015	Gorgan	type 2	330	51	WHOQOL		
[91]	Bahadori-Khosroshahi J	2011	Tabriz		100	20-60	WHOQOL-26	47.48	16.33
[92]	Fooladvandi M	2014	Kerman	type 2	96	53.08	SF-36	54.21	15.16
[93]	Shahraki-Vahed A	2010	Zabol	typ 1 and type 2	100	>7	SF-36		
[94]	Taghdisi MH	2011	Minudasht	type 2	78	49	WHOQOL-BREF	80.39	11.35
[95]	Sepehrnia I	2011	Karaj		30	40-65	SF-36	53.97	13.09
[96]	Fathi-Ahmadsaraee N	2016	Karaj	type 2	40	42.83	DQOL	26.37	4.51
[97]	Moein M	2014	Kashan	type 2	96	51.45	DQOL	105.23	16.06
[98]	Khalili M	2016	Isfahan	type 2	123	52	DQOL	1.88	0.36
[99]	Hadipour M	2013		type 2	3472	59.4	EQ-5D	0.623	0.387
[100]	Daneshvar S	2018	Ilam	typ 1 and type 2	122	57.74	SF-36		
[101]	Soleimani Z	2016	Sabzevar	typ 1 and type 2	189	51.7	DQOL-BCI	3.4	0.62
[102]	Kaveh MH	2018	Shiraz	type 2	207	55.35	DQOL	45.95	9.67
[103]	Shafiee-Kandjani AR	2018	Tabriz	type 2	263		SF-36	57.52	20.18
[104]	Sotodeh-Asl N	2020	Semnan	type 2	50	>18	SF-36	75.66	12.97
[105]	Tafazoli M	2017	Mashhad	type 2	90	43.58	SF-36	58.75	16.24
[106]	Tavakkoli L	2017	Kerman	type 2	198	54.91	WHOQOL-BREF		
[107]	Borhaninejad, VR	2016	Kerman		120	71.32	SF-36	46.48	20.45
[108]	Zareipour MA	2017		type 2	250	35-65	SF-36	58.32	19.62
[109]	Soleymanian T	2017	Tehran		219	62.2	SF-36	45.7	20.9
[110]	Barzegar Damadi MA	2018	Sari	type 2	15	43.5	D-39	151	33.17
[111]	Shakeri M	2018	Bojnord	type 2	18	53.5	SF-36		
[112]	Marzban A	2018	Yazd	type 2	600	56.11	DQOL	79.34	11.02
[113]	Ghaedrahmati A	2019	Isfahan	type 2	12	44	SF-36	54.25	4.78

Table 2: The mean QOL score in diabetic patients in Iran based on separate reviewed questionnaires

Questionnaire	Subgroups: Diabetic Patients' Quality of Life	Number of studies	The quality of life of diabetic patients (CI 95%)	P	I ² (%)
WHOQOL	Total	10	66.55 (45.83, 87.26)	<0.0001	100
	Men	4	46.41 (14.76, 78.06)	<0.0001	99.8
	Women	4	42.33 (13.92, 70.73)	<0.0001	99.9
	Physical Aspect	15	41.06 (26.35, 55.78)	<0.0001	100
	Mental Aspect	14	36.29 (22.26, 50.33)	<0.0001	100
	Social Activity Aspect	14	48.36 (34.63, 62.09)	<0.0001	100
	Peripheral Aspect	11	36.73 (29.46, 44)	<0.0001	99.9
	General Health Aspect	2	31.70 (-24.34, 87.73)	<0.0001	100
SF-36	Total	19	65.64 (59.82, 71.46)	<0.0001	98.3
	Men	1	49.86 (42.34, 57.38)	-	-
	Women	1	63.62 (56.64, 70.60)	-	-
	Physical Aspect	32	51.97 (42.75, 61.19)	<0.0001	100
	Mental Aspect	31	46.68 (38.99, 54.36)	<0.0001	99.9
	Social Activity Aspect	28	48.42 (41.37, 55.46)	<0.0001	99.9
	Peripheral Aspect	2	24.10 (22.94, 25.26)	0.143	53.4
	Vitality Aspect	24	49.69 (43.26, 56.11)	<0.0001	99.5
	General Health Aspect	24	43.62 (37.0, 50.24)	<0.0001	99.6
	Physical Pain Aspect	26	51.16 (40.61, 61.70)	<0.0001	99.9
	Physical Role Aspect	12	48.31 (42.53, 54.10)	<0.0001	96.5
	Emotion Aspect	15	51.32 (45.18, 57.47)	<0.0001	98.7
	Limitation of Activity Aspect	12	52.72 (33.13, 72.31)	<0.0001	99.7
	SF-20	Total	2	46.50 (37.19, 55.81)	0.002
Men		1	54.80 (49.87, 59.73)	-	-
Women		1	48.47 (44.71, 52.23)	-	-
Physical Aspect		1	16.05 (15.42, 16.68)	-	-
Mental Aspect		1	20.75 (19.99, 21.51)	-	-
Social Activity Aspect		1	18.05 (17.43, 18.67)	-	-
Peripheral Aspect		1	9.60 (9.06, 10.14)	-	-
DQOL	Total	10	61.19 (35.73-86.66)	<0.0001	100
	Physical Aspect	3	19.81 (8.70, 30.92)	<0.0001	99.9
	Mental Aspect	3	23.67 (10.00, 37.34)	<0.0001	99.9
	Social Activity Aspect	3	13.46 (7.03, 19.89)	<0.0001	99.6
	Peripheral Aspect	2	15.26 (-0.92, 31.44)	<0.0001	99.9
	General Health Aspect	1	41.25 (37.54, 44.96)	-	-
QOL	Total	3	117.91 (-62.97-298.79)	<0.0001	100
	Physical Aspect	2	9.95 (-5.40, 25.29)	<0.0001	99.8
	Mental Aspect	2	8.84 (-3.54, 21.23)	<0.0001	99.7
	Social Activity Aspect	2	5.18 (2.09, 8.26)	<0.0001	98.8
	Peripheral Aspect	1	20.23 (19.13, 21.33)	-	-
PedsQL	Total	2	34.36 (-31.49, 100.22)	<0.0001	100
	Physical Aspect	2	35.06 (-31.78, 101.89)	<0.0001	99.9
	Mental Aspect	2	34.29 (-30.44, 99.03)	<0.0001	99.9
	Social Activity Aspect	2	38.62 (-36.12, 113.37)	<0.0001	99.9
	Peripheral Aspect	2	33.15 (-30.76, 97.07)	<0.0001	99.9
	Emotion Aspect	1	59.84 (55.71, 63.97)	-	-
ADDQOL	Total	2	41.76 (12.01, 71.50)	<0.0001	99.4
	Physical Aspect	1	-1.81 (-1.96, -1.66)	-	-
	Mental Aspect	1	-0.94 (-1.11, -0.76)	-	-
	Social Activity Aspect	1	-0.96 (-1.07, -0.85)	-	-
D-39	Total	2	129.43 (88.77, 170.10)	<0.0001	95.4

Contd...

Table 2: Contd...

Questionnaire	Subgroups: Diabetic Patients' Quality of Life	Number of studies	The quality of life of diabetic patients (CI 95%)	P	I ² (%)
SWED-QUAL	Total	3	59.19 (21.15, 97.23)	<0.0001	99.8
	Physical Aspect	2	21.84 (14.66, 29.02)	<0.0001	98.9
	Physical Pain Aspect	2	8.07 (3.89, 12.26)	<0.0001	98.8
	Physical Role Aspect	1	9.70 (8.89, 10.51)	-	-
	Emotion Aspect	2	20.48 (9.50, 31.47)	<0.0001	99.5
IRDQOL	Total	2	105.92 (102.73, 109.10)	0.952	0
	Physical Aspect	1	57.03 (56.65, 57.41)	-	-
	Mental Aspect	1	59.54 (59.29, 59.79)	-	-
	Social Activity Aspect	1	69.53 (69.16, 69.90)	-	-
PHG-2	Total	2	22.63 (-2.38, 47.64)	<0.0001	99.9
	Physical Aspect	1	16.43 (15.60, 17.26)	-	-
	Social Activity Aspect	1	16.04 (15.30, 16.78)	-	-
	Emotion Aspect	1	9.84 (9.13, 10.55)	-	-
IDQOL-BCI	Total	1	3.40 (3.31, 3.49)	-	-
	Mental Aspect	2	11.82 (11.36, 12.29)	0.690	0
	Social Activity Aspect	2	11.82 (11.36, 12.29)	0.690	0

Table 3: The QOL of diabetic patients in Iran in three levels (good, fair, and poor)

Questionnaire	Subgroups	Number of study	The QOL in diabetic patients (95%CI)	P	I ² (%)
SF-36	Good	3	15 (-2, 32)	<0.0001	100
	Fair	3	68 (53, 83)	<0.0001	100
	Poor	3	46 (0, 92)	<0.0001	100
SF-20	Good	3	29 (14, 44)	<0.0001	100
	Fair	3	35 (30, 39)	<0.0001	99.5
	Poor	3	36 (32, 41)	<0.0001	99.5
QOL	Good	2	36 (24, 47)	<0.0001	100
	Fair	1	29 (29, 30)	-	-
	Poor	2	45 (20, 71)	<0.0001	100
WHOQOL	Good	1	55 (55, 55)	-	-
	Fair	1	56 (55, 56)	-	-
	Poor	1	37 (37, 37)	-	-
SWED-QUAL	Good	2	62 (19, 105)	<0.0001	100
	Poor	2	38 (-5, 81)	<0.0001	100
IRDQOL	Good	1	11 (11, 11)	-	-
	Fair	1	23 (22, 23)	-	-
	Poor	1	66 (66, 66)	-	-

in QOL, 129.43 in D-39, 34.36 in PedsQL, 41.76 in ADDQOL7, 22.63 in IDQOL-BCI8, 3.40 in DQOL-BCI, 0.62 in EQ-5D, 61.00 in PHG-2, 105.92 in IRDQOL, 59.19 in SWED-QUAL.

So far, several meta-analyses have been conducted on the status of QOL in diabetic patients in Iran, which we will examine below: In a meta-analysis of T. Schram *et al.* (2009)^[114] in The Netherlands, the aim was to investigate the relationship between depression and quality of life in diabetic patients. All studies suggest a negative association between depressive symptoms and at least

one aspect of QOL in people with diabetes. People with diabetes with depressive symptoms also had a much lower QOL than diabetes.

1. In meta-analysis of Kiadaliri *et al.* (2013)^[115] 46 studies found that people with diabetes were less likely to have health-related quality of life (HRQoL) without diabetes. The study covered 20 of Iran's 30 provinces. Of these 46 studies, 5 were type 1 diabetes and 23 were type 2 diabetes, and other studies were a combination of different types of diabetes. However, our study covered the studies published until 2017, and therefore the number of studies studied in our study is about twice that of the 2013 meta-analysis. In 2016, Soleimannejad *et al.*^[20] Studied the QOL of diabetic patients in 10 studies. And we decided

7 Audit of Diabetes Dependent Quality of Life

8 Iranian version of the Diabetes Quality of Life Brief Clinical Inventory




Figure 2: The relationship between quality of life score in diabetic patients and the sample size

to update this study: In the previous meta-analysis the number of studies studied was 10, whereas in the present study 82 studies were reviewed.

- In previous meta-analysis, only studies using questionnaires SF-36 and SF-20 were evaluated. However, in the present meta-analysis, all available questionnaires (WHOQOL-BREF, SF-36, SF-20, DQOL, QOL, PedsQL, ADDQoL, Youth Diabetes QOL and IDQOL-BCI) have been reviewed and no restrictions have been imposed on the questionnaire.
- The number of samples studied in the previous meta-analysis was 1,082, while in the present study 15,571 diabetic patients were evaluated.
- In the present meta-analysis, the QOL score of diabetic patients was examined by type of questionnaire and by dimensions of questionnaires and compared with each other, whereas this was not the case in previous meta-analysis.
- Current meta-analysis covers studies published as of December 31, 2016, while previous meta-analysis has carried out resource search for year 2015
- Current meta-analysis, in addition to the databases used by the previous meta-analysis, it has also examined the Cochrane, Embase, and Medline databases. Given the above, the present study is more complete than the previous meta-analysis study.
- In the present meta-analysis, the QOL of diabetic patients was evaluated in the form of levels: Good, Moderate, and Poor. This issue was not presented in previous meta-analysis.

Recently, two meta-analysis has been published in this regard, which we refer to: In meta-analysis Mokhtari *et al.* (2018)^[116] of 5,472 samples, the mean physical dimension score in patients with type 2 diabetes (53.5, 95% CI: 43.1–63.9) and the mean mental dimension score (54.5, 95% CI: 47–61.9) was less. As the age of the samples increased, the mean HRQoL score in diabetic patients in Iran decreased significantly.

In a meta-analysis of Dehvan *et al.* 2019,^[117] the QOL of type 2 diabetes patients in Iran was examined. The mean QOL of patients with type 2 diabetes was 61.90 (95% CI: 54.40–69.40). The highest and lowest QOL was achieved in terms of social support (49.19) and mental health (42.96). In this study, the WHOQOL-BREF questionnaire was used to assess the QOL of diabetic patients and therefore the number of studies studied was limited (16 studies). However, in our study, we did not have any restrictions on the type of diabetes or the type of questionnaire. A meta-analysis of Khunkaew *et al.* (2018),^[118] 12 studies in Australia found this conclusion. Overall, the HRQOL of participants in the studies was poor on four of eight subscales in the SF-36: Physical functioning (42.75); role physical (20.61); general health (39.52); and vitality (45.73). The results of this study are almost consistent with the results of the present meta-analysis.^[113–116]

Thommasen *et al.* conducted a study on the people of China, Malaysia, and India. In China, the mean scores of physical functioning was 83.3, public health was 69.3, social functioning was 83.9, and mental health was 72.9. In Malaysia, the mean scores of physical functioning was 86.6, public health was 68.6, social functioning was 78.8, and mental health was 75. In India, the mean scores of physical functioning was 73.9, public health was 70.1, social functioning was 86.1, and mental health was 71.5.^[119]

QOL^[1,118,119] Given that varied data have been archived for QOL of diabetic patients, the present meta-analysis was used to obtain an accurate estimate of the QOL of diabetic patients.

Conclusions

In this study, the QOL of diabetic patients was evaluated according to different types of studied questionnaires. We found that QOL of diabetic patients was lower than normal society. According to the results, the highest and lowest mean QOL score in diabetic patients in Iran were related to the D-39 questionnaire (129.43) and the EQ-5D questionnaires (0.62), respectively.

Limitations of the study

The limitations of the present study include lack of access to the full text of articles, lack of sufficient data in some articles, lack of reference to mean and standard deviation of QOL score in diabetic patients in some studies, and lack of uniform distribution of studies in different regions of Iran.

Authors' contribution

MF, MR, MA and DS searched the literature and analyzed the papers. The extraction stage was performed by MR, MA and DS. DS, MF, AHD prepared the manuscript. All authors read and signed the final paper.

Supplement

S1: Abbreviated table

Full name	Abbreviated name
Quality of life	QOL
WHO Quality of Life-BREF	WHOQOL-BREF
Short Form-36	SF-36
Short Form-20	SF-20
Diabetes Quality of Life	DQOL
Quality of Life	QOL
The World Health Organization Quality of Life	WHOQOL
Pediatric Quality of Life Inventory	PedsQL
Audit of Diabetes Dependent Quality of Life	ADDQoL
Iranian version of the Diabetes Quality of Life Brief Clinical Inventory	IDQOL-BCI

Ethical considerations

Ethical issues (including plagiarism, data fabrication, double publication) have been completely observed by the authors.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Received: 05 Sep 19 Accepted: 20 Sep 20

Published: 15 May 21

References

- Darvish Poor Kakhki A, Saeedi J, Yaghmaie F, Majd H, Montazeri A. Quality of life of diabetic patients referred to Tehran hospitals in 2004. *Int J Endocrinol Metab* 2006;8:49-56.
- Sayari A, Gari D, Asadai Lari M. Assessing quality of life, global experiences and the need for action in Iran. *Teb va Tazkiye* 2001;30-5.
- Cote I, Gregoire J, Moisan J, Chabot I. Quality of life in hypertension: The SF-12 compared to the SF-36. *Can J Clin Pharmacol* 2004;11:232-8.
- Masoudi-Alavi N, Ghofranipor F, Ahmadi F, Rajab A, Babai G. Quality of life in diabetic patient refers to diabetic association of Iran *Behbood Journal* 2004;8:47-56.
- Leininger M. Quality of life from a transcultural nursing perspective. *Nurs Sci Q* 1994;7:22-8.
- Karolson I, Berglin E, Larson P. Sense of coherence: Qol before and after coronary artery bypass surgery longitudinal study. *J Aadv Nurs* 2000;31:1383-92.
- Monica S, Clare M, Lesley B. Diabetic foot care: Assessing the impact of care on the whole patient. *Practical Diabetes Int* 2000;17:147-51.
- Glasgow R, Ruggiero L, Eakin E, Dryfoos J, Chobanian I. Quality of life associated characteristics in a large national sample of adults with diabetes. *Diabetes Care* 1997;20:562-7.
- Kotsanos J, Marrfero D, Viginati J, Mathias A, Huster W, Boggs M, *et al.* Health related quality of life results from multinational clinical trial of insulin lispro. *Diabetes Care* 1997;20:948-58.
- Andrew J. Guideline and performance measures for diabetes. *Am J Manag Care* 2007;13(Suppl 2):S41-6.
- Gambert S. Are we up to the challenge? *Clin Geriatr* 2002;10:14-7.
- Wild S, Ruglie G, Gron A, Sicree R, King H. Global prevalence of diabetes: Estimates for the Year 2000 and projections for 2030. *Diabetes Care* 2004;27:1047-53.
- Lustman PJ, Anderson RJ, Freedland KE, de Groot M, Carney RM, Clouse RE. Depression and poor glycemic control: A meta-analytic review of the literature. *Diabetes Care* 2000;23:934-42.
- Michael J. Diabetes foundation, clinical diabetes. *Am Diabetes Assoc* 2008;26:77-82.
- Graham JE, Stoebner-May DG, Ostir GV, Snih SA, Peek MK, Markides K, *et al.* Health related quality of life in older Mexican Americans with diabetes. *Health Qual Life Outcomes* 2007;5:39.
- Abdel Gawad E. Quality of life in Saudis with diabetes. *Saudi J Disabil* 2002;8:163-8.
- Heine J. Diabetes in the next century: Challenges and opportunities. *Neth J Med* 1999;55:265-70.
- Sabiha I. WHO study group. Prevention of diabetes mellitus: Report of a WHO study group. *Pak Dev Rev* 1995;34:91.
- Azizi F, Gouya MM, Vazirian P, Dolatshahi P, Habibian S. Screening for type 2 diabetes in the Iranian national programme: A preliminary report la revue de sant' de la me'diterrane'e orientale. *East Mediterr Health J* 2003;9:1122-7.
- Soleimannejad K, Sarokhani D, Sarokhani M, Sayehmiri K, Ahmadi N. Quality of life in diabetic patients in Iran: A systematic review and meta-analysis method. *Int J Pharm Tech* 2016;8:21608-18.
- Von Elm E, Altman DG, Egger M, Pocock SJ, Götzsche PC, Vandembroucke JP, *et al.* The Strengthening the reporting of observational studies in epidemiology (STROBE) statement: Guidelines for reporting observational studies. *Prev Med* 2007;45:247-51.
- Aghamollaei T, Eftekhari H, Shojaezadeh D, Mohammad K, Nakhjavani M, Pour F. Behavior, metabolic control and health-related quality of life in diabetic patients at Bandar Abbas diabetic clinic. *Iranian J Publ Health* 2003;32:54-9.
- Aghamolaei T, Eftekhari H, Mohammad K, Sobhani A, Shojaezadeh D, Nakhjavani M, *et al.* Influence of educational intervention using interaction approach on behavior change, hemoglobin A1C and health-related quality of life in diabetic patients. *J School Pub Health Int Pub Health Res* 2005;3:1-2.
- Sadeghie-Ahari S, Arshi S, Iranpour M, Amani F, Siahpush H. The effect of complications of type 2 diabetes in quality of life in diabetic patients. *J Ardabil Univ Med Sci* 2008;8:394-402.
- Ahmadi A, Hasanzadeh J, Rahimi M, Laskari L. Factors affecting quality of life in type 2 diabetics in Chaharmahal va Bakhtiari province. *J North Khorasan Univ Med Sci* 2011;3:7-11.
- Alavi A, Parvin N, Salehian T, Samipour V. Comparing quality of life of children and adolescents with diabetes mellitus and healthy group from patient and parents perspective in Shahrekord. *J Kordestan Univ Med Sci* 2010;15:46-52.
- Baghianimoghadam M, Afkhami-Ardakani M. The effect of educational intervention on quality of life of diabetic patients type 2, referee to diabetic research centre of Yazd. *Horizon Med Sci* 2008;13:21-8.
- Bazzazian S, Besharat M, Ehsan H, Rajab A. The moderating role of coping strategies in relationship between illness perception, quality of life and HbA1c in patients with type I diabetes. *Iranian J Endocrinol Metab* 2010;12:213-21.
- Borzou S, Salavati M, Safari M, Hadadinejad S, Zandieh M, Torkaman B. Quality of life in type II diabetic patients referred to

- Sina Hospital, Hamadan. *Zahedan J Res Med Sci* 2011;13:42-6.
30. Safarabadi-Farahani T, Ali-Akbar M, Safarabadi-Farahani A, Haghani H. Quality of life in young people with type 1 diabetes in relation to age and gender. *Iran J Nurs* 2011;23:73-9.
 31. Ghanbari A, Kazemnezhad E. A comparative study on the quality of life not insulin dependent diabetic melitus (niddm) patients of the razi hospital of rasht. *Modares J Med Sci* 2004;7:69-80.
 32. Ghanbari A, Yekta Z, Roushan Z, Lakeh N. Assessment of factors affecting quality of life in diabetic patients in Iran. *Public Health Nurs* 2005;22:311-22.
 33. Haririan H, Moghadasian S, Aghajanlou A. Quality of life and its dimensions in diabetic patients referring to diabetes center, Tabriz university of medical sciences, 2007. *J Diabetes Metab Disord* 2009;9:152-60.
 34. Heydari M, Alhani F, Kazemnejad A, Moezi F. The effect of empowerment model on quality of life of diabetic adolescents. *Iran J Pediatr* 2007;17:87-94.
 35. Jafari P, Forouzandeh E, Bagheri Z, Karamizadeh Z, Shalileh K. Health related quality of life of Iranian children with type 1 diabetes: Reliability and validity of the Persian version of the PedsQL™ generic core scales and diabetes module. *Health Qual Life Out* 2011;9:104.
 36. Ghavami H, Ahmadi F, Entezami H, Memarian R. The effect of continuous care model on quality of life in diabetic patients. *J Urmia Univ Med Sci* 2005;16:22-7.
 37. Jahanlou AS, Karami NA. The effect of literacy level on health related-quality of life, self-efficacy and self-management behaviors in diabetic patients. *Acta Med Iran* 2011;49:153-8.
 38. Jahanlou AS, Ghofranipour F, Kimmiagar M, Vafaei M, Heydarnia A, Sobhani A. Can quality of life questionnaires be used in diabetics to assess the relation between HbA1c and patients' domain aspects? *Acta Med Iran* 2011;49:246-51.
 39. Sedaghti-Kasbakh M, Ehsani M, Ghanbari A. Comparison of quality of life in patients with type 2 diabetes with non-diabetic patients. *J Babol Univ Med Sci* 2007;9:55-60.
 40. Kermansaravi F, Navidian A, Ansarymoghdamn A. Quality of life in type 1 diabetic adolescents in Zahedan (2011). *Iranian J Endocrinol Metab* 2012;13:651-7.
 41. Khaleidi S, Moridi G, Gharibi F. Survey of eight dimensions quality of life for patients with diabetes type II, referred to Sanandaj diabetes center in 2009. *J Fasa Univ Med Sci* 2011;1:29-37.
 42. Khamseh M, Monavari A, Malek M, Shafiee G, Baradaran H. Health-related quality of life in patients with type 1 diabetes. *Iranian J Endocrinol Metab* 2011;13:249-56.
 43. Peymani M, Monjamed Z, Aliasghapour M, Mehran A. Surveying the quality of life of diabetic patients with cardiovascular complications referring to the endocrinology clinic of selected hospitals of Tehran university of medical sciences in 2004-2005. *J Med Council IRI* 2007;25:142-8.
 44. Rakhshanderu S, Heydarnia A, Rajab A. The effect of health education on quality of life in diabetic patients. *Daneshvar Med* 2006;13:15-20.
 45. Rasouli D, Nasiriziba F, Nabiamjad R, Haghani H. Comparison of life quality in men and women with diabetic foot ulcer in selected hospitals of Tehran universities, 1387. *Pars J Med Sci* 2011;9:38-45.
 46. Safavi M, Samadi N, Mahmoodi M. Effect of quality of life improvement on type 2 diabetes patients' self-esteem. *Saudi Med J* 2011;32:954-7.
 47. Sanjari M, Safari S, Shokoohi M, Safizade H, Rashidinezhad H, Mashrouteh M, *et al.* A cross-sectional study in Kerman, Iran, on the effect of diabetic foot Ulcer on health-related quality of life. *Int J Low Extrem Wounds* 2011;10:200-2006.
 48. Shahrjerdi S, Shavandi N, Golpaygani M, Sheykhoseini R. Effect of endurance and strength training on blood glucose control, quality of life and mental health in women with type 2 diabetes. *Iranian J Diabetes Metab* 2009;9:35-44.
 49. Sayadi N, Fayazi S, Ramazani A. Comparison of quality of life among diabetic and non diabetic patients after open heart surgery (Short report). *J Rafsanjan Univ Med Sci Health Serv* 2011;10:144-50.
 50. Taghdisi M, Borhani M, Solhi M, Afkari M, Hosseini F. The effect of an education program utilising PRECEDE model on the quality of life in patients with type 2 diabetes. *Health Educ J* 2012;71:229-38.
 51. Timareh M, Rahimi M, Abbasi P, Rezaei M, Hyaidarpoor S. Quality of life in diabetic patients referred to the diabete research center in Kermanshah. *J Kermanshah Univ Med Sci (BEHBOOD)* 2012;16:63-9.
 52. Vares Z, Zandi M, Baghaei P, Alavi N, Ajourpaz N. Quality of life and related factors in diabetic patients attending a diabetes center in Kashan. *Nurs Res* 2010;5:14-22.
 53. Vazirinejad R, Sajadi MA, Maghool N. A historical cohort study assessing the effect of diabetes on the quality of life of patients. *Res Med* 2010;34:35-40.
 54. Yekta Z, Pourali R, Ghasemi-Rad M. Comparison of demographic and clinical characteristics influencing health-related quality of life in patients with diabetic foot ulcers and those without foot ulcers. *Diabetes Metab Syndr Obes* 2011;4:393-9.
 55. Mirfeizi M, Jafarabadi M, Toorzani Z, Mohammadi S, Azad M, Mohammadi A, *et al.* Feasibility, reliability and validity of the Iranian version of the diabetes quality of life brief clinical inventory (IDQOL-BCI). *Diabetes Res Clin Pract* 2012;96:237-47.
 56. Shahi M, Mohammadyfar M. Comparison of depression, anxiety, stress, quality of life, and alexithymia between people with type II diabetes and non-diabetic counterparts. *Pers Individ Differ* 2017;104:64-8.
 57. Najafi-Ghezjeljeh T, Kohandany M, Oskouei F, Malek M. The effect of progressive muscle relaxation on glycated hemoglobin and health-related quality of life in patients with type 2 diabetes mellitus. *Appl Nurs Res* 2017 33:142-8.
 58. Shamshirgaran S, Ataei J, Alamdari M, Safaeian A, Aminisani N. Predictors of health-related quality of life among people with type II diabetes Mellitus in Ardabil, Northwest of Iran, 2014. *Prim Care Diabetes* 2016 10:244-50.
 59. Hajian-Tilaki K, Heidari B, Hajian-Tilaki A. Solitary and combined negative influences of diabetes, obesity and hypertension on health-related quality of life of elderly individuals: A population-based cross-sectional study. *Diabetes Metab Syndr* 2016;10:37-42.
 60. Dadgostar H, Firouzinezhad S, Ansari M, Younespour S, Mahmoudpour A, Khamseh M. Supervised group-exercise therapy versus home-based exercise therapy: Their effects on quality of life and cardiovascular risk factors in women with type 2 diabetes. *Diabetes Metab Syndr* 2016;10:30-6.
 61. Jafari N, Farajzadegan Z, Loghmani A, Majlesi M, Jafari N. Spiritual well-being and quality of life of Iranian adults with type 2 diabetes. *Evid Based Complement Alternat Med* 2014;2014:8.
 62. Abdoli S, Abdoli S. Quality of life in people with type 2 diabetes living in rural and remote areas, Iran. *Int J Diabetes Dev Ctries* 2015;35:290-7.
 63. Hadi N, Ghahramani S, Montazeri A. Health related quality of life in both types of diabetes in Shiraz, Iran. *Shiraz E Medical J*

- 2013;14:112-22.
64. Shavandi N, Sharjerdi S, Sheikh Hoseini R, Ghorbani A. The effect of strengthening exercises on metabolic factors, quality of life and mental health in women with type 2 diabetes. *Iranian J Endocrinol Metab* 2010;12:222-30.
 65. Shayeghian Z, Aguilar-Vafaie M, Besharat M, Parvin M, Roohi Gilani K. The association between self-care and control of blood sugar and health-related quality of life in type II diabetes patients. *Iranian J Endocrinol Metab* 2014;15:545-51.
 66. Alipour A, Zare H, Poursharifi H, Sheibani K, Ardekani M. The intermediary role of self-efficacy in relation with stress, glycosylated haemoglobin and health-related quality of life in patients with type 2 diabetes. *Iran J Public Health* 2012;41:76-80.
 67. Afshar M, Memarian R, Mohammadi E. The effect of group discussion on the quality of life and HbA1c levels of adolescents with diabetes. *Iran Red Crescent Med J* 2014;16:e21110.
 68. Derakhshanpour F, Vakili M, Farsinia M, Mirkarimi K. Depression and quality of life in patients with type 2 diabetes. *Iran Red Crescent Med J* 2015;17:e27676.
 69. Zaker M, Moghadam A, Shams S, Moradi Y. Effect of self-management education package on specific quality of life among diabetic patients in Urmia diabetes centers. *J Chem Pharm Sci* 2016;9:1396-9.
 70. Didarloo A, Alizadeh M. Health-related quality of life and its determinants among women with diabetes mellitus: A cross-sectional analysis. *Nurs Midwifery Stud* 2016;5:e28937.
 71. Gholami A, Azini M, Borji A, Shirazi F, Sharafi Z, Zarei E. Quality of life in patients with type 2 diabetes: Application of WHOQOL-BREF scale. *Shiraz E Medical J* 2013;14:162-71.
 72. Torabi M, Izadi A, Naderi F, Shamsaei F. Sleep quality and quality of life in adults with type 2 diabetes. *J Diabetes Nurs* 2014;2:51-61.
 73. Izadi A, Sepahvand F, Naderifar M, Mohammadipour F. The effect of an educational intervention on quality of life in patients with type 2 diabetes referring to Tamin Ejtemaei hospital in 2013. *J Diabetes Nurs* 2014;2:18-28.
 74. Khodabakhshi-Koolaei A, Navidian A, Baiati Z, Rahmatizadeh M. Effectiveness of supportive psychotherapy on quality of life in patients with type 2 diabetes. *J Diabetes Nurs* 2015;3:31-41.
 75. Mohammadshahi M, Shirani F, Elahi S, Ghasemi S, Shahni M, Haidari F. Evaluation of relationship between dietary patterns and quality of life in patients with type 2 diabetes. *Daneshvar Med* 2015;22:1-12.
 76. Saeedpour J, Jafari M, Asgar M, Dardashti H, Teymoozadeh E. The impact of self-care education on life quality of diabetic patients. *J Health Admin* 2013;16:26-36.
 77. Ghasemipour M, Ghasemi V, Zamani A. Evaluation of quality of life and its dimensions in diabetic patients referring to Khorramabad Shohada hospital (in 2008). *Yafteh* 2009;11:125-33.
 78. EydiBaygi M, Mehrabizade M, Davoudi I, Ahmadi V, Dehghanizade Z. Comparison the quality of life in patients with diabetes type 2 and non-diabetic individuals. *Sci J Ilam Uni Med Sci* 2014;22:55-62.
 79. Sadeghi T, Derakhshan R. The effect of nurse telephone follow up on quality of life in diabetic patients. *Payesh* 2012;11:711-7.
 80. Zaree-Bahramabadi M, Vafaei-Baneh F, Ghaderi E, Taghvaei D. The Effectiveness of cognitive-behavioral therapy on quality of life in patients with type 2 diabetes. *J Diabetes Metab Disord* 2013;12:225-32.
 81. Ghashghaie S, Farnam R. The effectiveness of mindfulness-based cognitive therapy on quality-of-life in outpatients with diabetes. *Iranian J Diabetes Metab* 2014;13:319-30.
 82. Saadatjoo S, Rezvanee M, Tabyee S, Oudi D. Life quality comparison in type 2 diabetic patients and none diabetic persons. *Mod Care Sci Qlty Birjand Nurs Midwifery Fac* 2012;9:24-31.
 83. Behrouz B, Bavali F, Heidarizadeh N, Farhadi M. The effectiveness of acceptance and commitment therapy on psychological symptoms, coping styles, and quality of life in patients with type-2 diabetes. *J Health* 2016;7:236-53.
 84. Ebrahimi H, Sadeghi M, Bazghaleh M, Shaker S, Ghasemi M. Relationship between metabolic control indexes and quality of life in patients with type II diabetes mellitus. *Iran J Nurs* 2014;27:73-82.
 85. Mohammad Rahimi G, Attarzadeh Hosseini S. Effect of aerobic training and diet on insulin resistance and quality of life in type II diabetic patients. *Horizon Med Sci* 2016;21:277-84.
 86. Shams S, Zaker M, Ghavami H. Effect of self management educative package on quality of life among diabetic patients in Urmia diabetes centers between in the years of 2013. *J Urmia Nurs Midwifery Fac* 2016;13:863-8.
 87. Mohammadpour Y, Haririan H, Moghadasian S, Ebrahimi H. Study of quality of life and its dimensions in diabetic patients referring to diabetes center of Tabriz university of medical sciences in 2007. *J Urmia Nurs Midwifery Fac* 2008;6:26-37.
 88. Ganjlo J, Talebi Z, Assarroudi A, Rakhshani M. Comparative assessment of effect of education in the OREM's self care model way with current method on the quality of life of diabetic type 2 patients. *Bimonthly J Sabzevar Univ Med Sci* 2015;22:748-57.
 89. Bidi F, Hassanpour K, Ranjbarzadeh A, Kheradmand A. Effectiveness of educational program on knowledge, attitude, self care and life style in patients with type II diabetes. *J Sabzevar Univ Med Sci* 2013;19:336-44.
 90. Derakhshanpoor F, Farsinia M, Shahini N. Relationship between anxiety disorders and life quality in type two diabetic patients. *J Res Dev Nurs Midw* 2015;12:94-102.
 91. Bahadori-Khosroshahi J, Khanjani Z. Depression and quality of life between diabetic and non-diabetic patients. *J Health Psychol* 2011;1:61-77.
 92. Fuladvandi M, Aziz Zadeh Foroozi M, Asad Abadi A, Fuladvandi G, Lashkari T, Malekian L. Effectiveness of stress management training on improved quality of life in patients with type 2 diabetes. *J Health Promot Manag* 2014;3:16-24.
 93. Shahraki-Vahed A, Hamed-Shahraki S, Masinaeinezhad N, Shahdadi H. Survey of life's quality in diabetic patients referred to diabet's clinic of Zabol 2010. *J Rostamineh* 2011;3:21-9.
 94. Taghdisi M, Borhani M, Solhi M, Afkari M, Hosseini M. Effect of educational program based on PRECED model on quality of life in patients with typeII diabetes. *J Gorgan Univ Med Sci* 2011;13:29-36.
 95. Sepehrnia E, Ranjbar F, Hosseinzadeh-Taghvaei M, Peymani J. The effectiveness of life skills training on the subscale of quality of life in people with diabetes. *J Health Psychol* 2011;1:81-102.
 96. Fathi Ahmadsaraei N, Neshat Doost H, Manshaee G, Nadi M. The effectiveness of acceptance and commitment therapy on quality of life among patients with type 2 diabetes. *Iran J Health Educ Health Promot* 2016;4:31-9.
 97. Moein M, Aghajani M, Mirbagher-Ajorpaz N. Effects of the empowerment program on the quality of life in patients with type II diabetes. *J Diabetes Nurs* 2015;3:29-41.
 98. Khalili M, Sabouhi F, Abazari P, Aminorroaya A. Comparing the quality of life in insulin recipient and refusing patients with type 2 diabetes. *Iran J Nurs Midwifery Res* 2016;21:351-6.
 99. Hadipour M, Aboalhasani F, Molavi-Vardanjanji H. Health related quality of life in patients with of type II diabetes in Iran. *Payesh*

- 2013;12:135-41.
100. Daneshvar S, Khodamoradi A, Ghazanfari Z, Montazeri A. Quality of life in diabetic patients: A comparative study. *Payesh* 2018;17:541-50.
 101. Solimani Z, Barati H, Mozafari J, Ershadi M, Mohammadi M. The quality of life of patients with diabetes from the city of Sabzevar during year 2016. *Military Caring Sci* 2017;3:264-71.
 102. Kaveh M, Ghahremani L, Nazari M, Zare S. Quality of life in diabetic patients: The predicting role of personal resources. *J Health Sci Surveillance Sys* 2018;6:142-8.
 103. Shafee-Kandjani AR, Hosseinpour M, Shoja H, Daneshamouz H, Mohammad-Alizadeh S. Assessing quality of life and general health status in married women with type II diabetes in Tabriz. *Depiction Health* 218;9:195-205.
 104. Sotodeh Asl N, Avazabadian M, Ghorbani R, Malek F. Quality of life in patients with hypertension and type 2 diabetes mellitus. *Koumesh* 2020;22:263-8.
 105. Tafazoli M, Parnan A, Azmoude E. Sexual function and quality of life in diabetic women referring to health care centers in Mashhad. *J Edu Health Promot* 2017;6:25.
 106. Tavakkoli L, Dehghan A. Compare the quality of life in type 2 diabetic patients with healthy individuals (Application of WHOQOL-BREF). *Zahedan J Res Med Sci* 2017;19:e5882.
 107. Borhaninejad V, Kazazi L, Haghi M, Chehrehnegar N. Quality of life and its related factors among elderly with diabetes. *Iran J Ageing* 2016;11:162-73.
 108. Zareipour M, Ghelichi Ghoghogh M, Mahdi-Akhgar M, Alinejad M, Akbari S. The quality of life in relationship with glycemic control in people with type 2 diabetes. *J Community Health Res* 2017;6:141-9.
 109. Soleymanian T, Kokabeh Z, Mahjoub A, Ramaghi R, Argani H. Clinical outcomes and quality of life in hemodialysis diabetic patients versus non-diabetics. *J Nephropathol* 2017;6:81-9.
 110. Barzegar Damadi M, Mirzaian B, Akha O, Hosseini S, Jadidi M. Effect of cognitive-behavioral group therapy on HbA1C, self-efficacy, depression, illness perception, and quality of life in patients with type II diabetes. *J Mazandaran Univ Med Sci* 2018;27:87-100.
 111. Shakeri M, Hatami M, Hasani J, Shakeri HS. The Effectiveness of Mindfulness-Based Stress Reduction (MBSR) on mental health and quality of life of the patients with type 2 diabetes. *JNKUMS* 2018;10:31-40.
 112. Marzban A. Relationship between spiritual health and quality of life in type II diabetic patients: A cross-sectional study in Yazd. *J Diabetes Nurs* 2018;6:641-52.
 113. Ghaedrahmati A, Jabalameli S. Effect of acceptance and commitment therapy on the quality of life and physical indices of patients with diabetes. *J Diabetes Nurs* 2019;7:915-28.
 114. Schram M, Baan C, Pouwer F. Depression and quality of life in patients with diabetes: A systematic review from the European depression in diabetes (EDID) research consortium. *Curr Diabetes Rev* 2009;5:112-9.
 115. Kiadaliri AA, Najafi B, Mirmalek-Sani M. Quality of life in people with diabetes: A systematic review of studies in Iran. *J Diabetes Metab Disord* 2013;12:54.
 116. Mokhtari Z, Gheshlagh RG, Kurdi A. Health-related quality of life in Iranian patients with type 2 diabetes: An updated meta-analysis. *Diabetes Metab Syndr* 2019;13:402-7.
 117. Dehvan F, Saeid DM, Dehkordi AH, Gheshlagh RG. Quality of life of Iranian patients with type 2 diabetes: A systematic review and meta-analysis. *Nurs Pract Today* 2019;6:167-75.
 118. Khunkaew S, Fernandez R, Sim J. Health-related quality of life among adults living with diabetic foot ulcers: A meta-analysis. *Qual Life Res* 2019;28:1413-27.
 119. Thommasen HV, Zhang W. Impact of chronic disease on quality of life in the Bella Coola valley. *Rural Remote Health* 2006;6:528.