

The Prevention of Cisplatin-Induced Nephrotoxicity: A General Consensus Statement of a Group of Oncologist-Hematologists, Adult and Pediatric Nephrologists, Radiation Oncologists, Clinical Pathologists, Clinical Pharmacologists, and Renal Physiologists on Cisplatin Therapy in Cancer Patients

Abstract

Backgrounds: Most of the cancer patients with solid tumor are subjected to chemotherapy with cisplatin (CP) in clinic. However, the most side effect of CP is nephrotoxicity, which limits the treatment. The aim of study was to develop a general consensus statement for CP therapy in clinic to limit the drug-induced nephrotoxicity. **Methods:** A total of 30 oncologist-hematologists, adult and pediatric nephrologists, radiation oncologists, clinical pathologist clinical pharmacologist, and renal physiologist participated in a workshop, and in order to reduce the incidence of CP-induced nephrotoxicity, a general consensus was developed. **Results:** The developed general consensus was focused on some items such as age, sex, female hormone, nonsteroidal anti-inflammatory drugs (NSAID), renin-angiotensin system inhibitor drugs, glomerular filtration rate, hydration methods, contrasts, antioxidants, dextrose, and magnesium. **Conclusion:** The agreement between participants for CP therapy in clinic was achieved, and this general consensus was announced to be implemented in the hospitals.

Keywords: Cisplatin, consensus, nephron, toxicity

Introduction

Cisplatin (CP) is the most frequently used drug for cancer treatment. Acute kidney injury, glomerular capillary endothelial injury, tubulointerstitial disease, and renal electrolyte disorders are the most frequent adverse effects of CP therapy, which are major causes of morbidity and mortality among cancer patients. Volume depletion, preexisting kidney disease, cardiovascular disease, diabetes, and concurrent use of nonsteroidal other nephrotoxic drugs such as nonsteroidal anti-inflammatory drugs (NSAID) are the major risk factors for CP-induced nephrotoxicity.

There are some risk factors associated with the incidence of CP-induced nephrotoxicity.^[1] In addition, many other clinical and experimental studies have explained the CP-induced nephrotoxicity risk factors, including age,^[2-4] gender,^[5-7] low glomerular filtration rate (GFR),^[8] sex hormone estrogen,^[9] dehydration,^[8]

hydration process during CP therapy with dextrose,^[10] NSAID, and contrast media.^[11] Special attentions also were made to antioxidant therapy to attenuate CP-induced nephrotoxicity in the laboratory.^[12-16] Based on new findings, the general consensus will be helpful to prevent nephrotoxicity during CP therapy.

Methods

A total of 30 oncologist-hematologists, adult and pediatrics' nephrologists, radiation oncologists, clinical pathologists, clinical pharmacologist, and renal physiologists participated in a workshop to develop a general consensus to reduce the incidence of CP-induced nephrotoxicity in Isfahan, Iran hospitals. The announcement for the workshop was performed 2 months earlier, and the participants were asked to review the related materials in the literature including their experiences to obtain the general consensus questions. Based on the suggestions, four main subjects related to

Farzaneh Ashrafi^{1,2},
Mojgan Mortazavi^{3,4},
Mehdi
Nematbakhsh^{1,5,6}

¹Water and Electrolytes Research Center, Isfahan University of Medical Sciences, Isfahan, Iran, ²Department of Internal Medicine, Oncology-Hematology Section, Isfahan University of Medical Sciences, Isfahan, Iran, ³Isfahan Kidney Diseases Research Center, Isfahan University of Medical Sciences, Isfahan, Iran, ⁴Department of Internal Medicine, Nephrology Section, Isfahan University of Medical Sciences, Isfahan, Iran, ⁵Department of Physiology, Isfahan University of Medical Sciences, Isfahan, Iran, ⁶IsfahanMN Institute of Basic and Applied Sciences Research, Isfahan, Iran

Address for correspondence:
Prof. Mehdi Nematbakhsh,
Water and Electrolytes Research
Center, Isfahan University of
Medical Sciences, Isfahan, Iran.
E-mail: nematbakhsh@med.mui.ac.ir

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.net

DOI:
10.4103/ijpvm.IJPVM_445_19

Quick Response Code:



This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Ashrafi F, Mortazavi M, Nematbakhsh M. The prevention of cisplatin-induced nephrotoxicity: A general consensus statement of a group of oncologist-hematologists, adult and pediatric nephrologists, radiation oncologists, clinical pathologists, clinical pharmacologists, and renal physiologists on cisplatin therapy in cancer patients. *Int J Prev Med* 2022;13:21.

Table 1: A general consensus statement on cisplatin therapy in cancer patients: the associated items and factors and principals

General consensus parts	Items	Principals and descriptions
Part one: General consideration factors	Age and gender	Principal 1: CP-induced nephrotoxicity is age and gender related. ^[2-8,22] The age >60 years has more risk for CP-induced nephrotoxicity
	RAS inhibitor drugs	Principal 2: The drugs like losartan and enalapril may increase the risk of CP-induced nephrotoxicity in female gender ^[5,23,24] due to expression difference of RAS receptors between female and male
Part two: Prohibition items	Female sex hormone estrogen	Principal 3: Based on some laboratory studies, high levels of estrogen along with CP may increase the risk of CP-induced nephrotoxicity ^[9,25-28]
	eGFR	Principal 4: Determining the eGFR before treatment with the CP is necessary and important ^[29,30]
	GFR >50 ml/min per 1.73 m ² body surface	Principal 5: Although when eGFR is less than 60 ml/min/1.73 m ² body surface, ^[8] it is not recommended for CP treatment; however, based on other finding ^[29] and our clinical experience, CP treatment is advised when the eGFR is greater than 50 ml/min/1.73 m ² body surfaces. It is important that when GFR is less than 50 ml/min per 1.73 m ² body surface, the administration of CP can be made only with the approval of a multidisciplinary consulting medical team
	Hydration with dextrose	Principal 6: Hydration with dextrose (dextrose diuresis) does not attenuate CP-induced nephrotoxicity and should be avoided because dextrose may increase the risk of CP-induced nephrotoxicity. ^[10,31] Before and after administration of CP, hydration should be performed with normal saline plus potassium chloride (KCl)
	Mannitol	Principal 7: There is controversy regarding the use of mannitol for diuresis. On the basis of many agreed and opposed studies related to hydration with mannitol, ^[17-21,32,33] special care is needed when mannitol administration is applied by a multidisciplinary consulting medical team
	NSID*	Principal 8: There is interaction between NSID and CP; therefore, using NSID with CP increases the risk of CP-induced nephrotoxicity and must be avoided ^[4,11]
	Contrast media*	Principal 9: Co-administration of nephrotoxic drug such as CP with contrast media is a high-risk factor to develop CP-induced nephrotoxicity ^[4,11]
Part three: Suitable hydration requirements and permissions items	Antioxidant supplements*	Principal 10: There are a lot of basic researches and less clinical research about the use of antioxidants to reduce CP-induced nephrotoxicity. ^[12-14, 34-36] However, prescription of herbal or synthetic antioxidants supplements to reduce CP-induced nephrotoxicity without clinical scientific citation or the approval of a multidisciplinary consulting medical team is not recommended
	Suitable hydration agents	Principal 11: Before and after administration of CP, hydration should be performed with normal saline (1 1/2-4 h) plus KCl
Part four: Others	Magnesium administration	Principal 12: There are numerous basic and clinical studies that make it doubtful about the role of magnesium in prevention of CP-induced nephrotoxicity; ^[37-45] therefore, administration of intravenous magnesium with hydration is recommended only if serum level of magnesium is lower than normal
	Diabetes, acidosis, hypokalemia, uricosuric, hyperuricemia, infection, etc.	Principal 13: There are many other pathological conditions such as diabetes, kidney transplantation, acidosis, hypokalemia, hyperuricemia, antibiotics, infection, etc. that need to be considered on a case-by-case basis

*Nephrotoxic antibiotic such as aminoglycoside, vancomycin, and amphotericin B that are commonly used to treat infections in patients with cancer. RAS: Renin-angiotensin system; eGFR: Estimated GFR; NSID: Nonsteroidal anti-inflammatory drugs; KCl: Potassium chloride

CP therapy were assigned for the workshop discussion as followings:

- (1) Based on the clinical evidences, to reduce the CP-induced nephrotoxicity, the CP prescribing should be accompanied by hydration of the patient. Therefore, the agreement on the hydration process seems necessary.
- (2) Based on the experimental evidences, some synthetic and herbal antioxidant supplements can reduce CP-induced nephrotoxicity, and it seems necessary to agree on this issue in the process of CP therapy.
- (3) From the clinical point of view, the patients should have favorable clinical conditions for receiving CP (e.g., the

minimum level of GFR) and therefore agreeing to these issues in the CP prescribing process seems essential.

- (4) Some drugs along with CP increases the risk of nephrotoxicity and it is therefore necessary to agree on these issues in the process of CP prescribing.

The subjects with the 16 questions were submitted to the participants before the workshop day. On the day of workshop, two nephrologists and two oncologist-hematologists also were asked to summarize the literature reviewing results. During the discussion, some other questions also were created by the participants.

Therefore, all questions, one by one, were subjected to panel discussion, and based on the opinion of all, the agreement was created. Finally, three workshop members were assigned to prepare the draft of general consensus, and then the draft was finalized. This general consensus included 13 principals or important items that must be considered during CP therapy to limit CP-induced nephrotoxicity.

Results

The general consensus included four different parts.

Part one: general consideration factors include 4 items (principals 1–4)

The general consideration factors were listed in Table 1. Age, gender, concurrent use of renin–angiotensin system inhibitor drugs, the level of female sex hormone estrogen, and estimated GFR should be considered in the first step.

Part two: prohibition items include 6 items (principals 5–10)

To avoid CP-induced nephrotoxicity, some items are forbidden during CP therapy, including GFR <50 ml/min/1.73 m² body surface, hydration with dextrose, NSAID prescribing, given contrast media during 2 weeks post-CP administration, and prescribing antioxidant without enough clinical evidences [Table 1]. Due to existence of positive and negative results for mannitol prescribing,^[17-21] special care is recommended when mannitol administration is applied.

Part three: requirement and permission items include 2 items (principals 11–12)

There are two major requirements and a major license. Before and after administration of CP, hydration should be performed with normal saline plus potassium chloride (KCl). Administration of intravenous magnesium with hydration is recommended only when the serum level of magnesium is lower than normal [Table 1].

Part four: Others include 1 item (principal 13)

There are many other pathological conditions such as diabetes, kidney transplantation, acidosis, hypokalemia, hyperuricemia infection, etc. that need to be considered on a case-by-case basis.

Conclusion

We suggested several new points that need to be considered during CP administration. Female sex hormone estrogen,^[9,25-28] hydration with dextrose,^[10,31] and concurrent use of renin–angiotensin system inhibitor drugs^[5,23,24] are the subjects that included in this general consensus. In order to obtain a more complete general consensus or guideline for CP therapy, more clinical trial researches are needed.

Author contribution

All the authors (FA, MM, and MN) were contributed to design the workshop, to search the main literature review, to manage the workshop, and to write the article draft. The article was finalized by MN.

Acknowledgements

Special thanks to the entire workshop participants for their thoughtful reviewing and discussing; Dr Fariborz Mokarian, Dr Shahrzad Shahidi, Dr Mohammad Matinfar, Dr Shahram Taheri, Dr Abdolamir Atapoor, Dr Hamid Nasri, Dr Valiollah Mahrzad, Dr Alaleh Gheisari, Dr Shiva Seirafian, Dr Mehdi Mahmoodzadeh, Dr Firoozeh Moeinzadeh, Dr Mehran Sharifi, Dr Alireza Sadeghi, Dr Nooshin Hashemi, Dr Maryam Mirpoorian, Dr Padideh Oghab, Dr Reihaneh Motamedifard, Dr Ali Akhavan, Dr Farhad Ghadiri, Dr Ali Derakhshandeh, Dr Yaser Salehi, Dr Mohammad-Reza Khosravi, Dr Shirinsadat Badri, Dr Ziba Farajzadeghan, Dr Ardeshir Talebi, Dr Yousef Gheisari, and Dr Aliraza Andalib. Additional thanks to Dr Fariborz Mokarian, Dr Shahrzad Shahidi, Dr Mohammad Matinfar for their lectures in the workshop.

Ethical approval

This article does not contain any studies with human participants or animals performed by any of the authors.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Received: 28 Nov 19 **Accepted:** 27 Nov 20

Published: 08 Feb 22

References

1. Sanchez-Gonzalez PD, Lopez-Hernandez FJ, Lopez-Novoa JM, Morales AI. An integrative view of the pathophysiological events leading to cisplatin nephrotoxicity. *Crit Rev Toxicol* 2011;41:803-21.
2. de Jongh FE, van Veen RN, Veltman SJ, de Wit R, van der Burg ME, van den Bent MJ, *et al.* Weekly high-dose cisplatin is a feasible treatment option: Analysis on prognostic factors for toxicity in 400 patients. *B J Cancer* 2003;88:1199-206.
3. Pezeshki Z, Maleki M, Talebi A, Nematbakhsh M. Age and gender related renal side effects of cisplatin in animal model. *Asian Pac J Cancer Prev* 2017;18:1703-5.
4. Ali BH, Al-Moundhri M, Tageldin M, Al Husseini IS, Mansour MA, Nemmar A, *et al.* Ontogenic aspects of cisplatin-induced nephrotoxicity in rats. *Food Chem Toxicol* 2008;46:3355-9.
5. Nematbakhsh M, Pezeshki Z, Eshraghi Jazi F, Mazaheri B, Moeini M, Safari T, *et al.* Cisplatin-induced nephrotoxicity; protective supplements and gender differences. *Asian Pac J Cancer Prev* 2017;18:295-314.
6. Aydin I, Agilli M, Aydin FN. Gender differences influence renal injury in cisplatin-treated rats: Biochemical evaluation. *Biol Trace Elem Res* 2014;158:275.

7. Nematbakhsh M, Ebrahimi S, Toyserkani M, Eshraghi-Jazi F, Talebi A, Ashrafi F. Gender difference in Cisplatin-induced nephrotoxicity in a rat model: Greater intensity of damage in male than female. *Nephrourol Mon* 2013;5:818-21.
8. Isnard-Bagnis C, Moulin B, Launay-Vacher V, Izzedine H, Tostivint I, Deray G. [Anticancer drug-induced nephrotoxicity]. *Nephrol Ther* 2005;1:101-14.
9. Pezeshki Z, Nematbakhsh M, Nasri H, Talebi A, Pilehvarian AA, Safari T, *et al.* Evidence against protective role of sex hormone estrogen in Cisplatin-induced nephrotoxicity in ovariectomized rat model. *Toxicol Int* 2013;20:43-7.
10. Karimi F, Kassaei S, Baradaran A, Ashrafi F, Talebi A, Lak Z, *et al.* Dextrose hydration may promote cisplatin-induced nephrotoxicity in rats: Gender-related difference. *Indones Biomed J* 2019;11:136-44.
11. Jones DP, Spunt SL, Green D, Springate JE, Children's Oncology G. Renal late effects in patients treated for cancer in childhood: A report from the Children's oncology group. *Pediatr Blood Cancer* 2008;51:724-31.
12. Kamel KM, Abd El-Raouf OM, Metwally SA, Abd El-Latif HA, El-sayed ME. Hesperidin and rutin, antioxidant citrus flavonoids, attenuate cisplatin-induced nephrotoxicity in rats. *J Biochem Mol Toxicol* 2014;28:312-9.
13. Somani SM, Husain K, Whitworth C, Trammell GL, Malafa M, Rybak LP. Dose-dependent protection by lipoic acid against cisplatin-induced nephrotoxicity in rats: Antioxidant defense system. *Pharmacol Toxicol* 2000;86:234-41.
14. Nematbakhsh M, Pezeshki Z. Sex-related difference in nitric oxide metabolites levels after nephroprotectant supplementation administration against cisplatin-induced nephrotoxicity in wistar rat model: The role of vitamin E, erythropoietin, or N-Acetylcysteine. *ISRN Nephrol* 2013;2013:612675.
15. Nematbakhsh M, Ashrafi F, Safari T, Talebi A, Nasri H, Mortazavi M, *et al.* Administration of vitamin E and losartan as prophylaxes in cisplatin-induced nephrotoxicity model in rats. *J Nephrol* 2012;25:410-7.
16. Atasayar S, Gurer-Orhan H, Orhan H, Gurel B, Girgin G, Ozgunes H. Preventive effect of aminoguanidine compared to vitamin E and C on cisplatin-induced nephrotoxicity in rats. *Exp Toxicol Pathol* 2009;61:23-32.
17. El Hamamsy M, Kamal N, Bazan NS, El Haddad M. Evaluation of the effect of acetazolamide versus mannitol on cisplatin-induced nephrotoxicity, a pilot study. *Int J Clin Pharm* 2018;40:1539-47.
18. Dhillon P, Amir E, Lo M, Kitchlu A, Chan C, Cochlin S, *et al.* A case-control study analyzing mannitol dosing for prevention of cisplatin-induced acute nephrotoxicity. *J Oncol Pharm Pract* 2019;25:875-83.
19. Adam JP, Fournier MA, Letarte N. Should mannitol be reinstated in prevention of cisplatin-induced nephrotoxicity? *J Oncol Pharm Pract* 2018;24:239-40.
20. Morgan KP, Buie LW, Savage SW. The role of mannitol as a nephroprotectant in patients receiving cisplatin therapy. *Ann Pharmacother* 2012;46:276-81.
21. Williams RP, Jr., Ferlas BW, Morales PC, Kurtzweil AJ. Mannitol for the prevention of cisplatin-induced nephrotoxicity: A retrospective comparison of hydration plus mannitol versus hydration alone in inpatient and outpatient regimens at a large academic medical center. *J Oncol Pharm Pract* 2017;23:422-8.
22. Launay-Vacher V, Isnard-Bagnis C, Janus N, Karie S, Deray G. [Chemotherapy and renal toxicity]. *Bull Cancer* 2008;95 FMC Onco:F96-103.
23. Zamani Z, Nematbakhsh M, Eshraghi-Jazi F, Talebi A, Jilanchi S, Navidi M, *et al.* Effect of enalapril in cisplatin-induced nephrotoxicity in rats; gender-related difference. *Adv Biomed Res* 2016;5:14.
24. Haghighi M, Nematbakhsh M, Talebi A, Nasri H, Ashrafi F, Roshanaei K, *et al.* The role of angiotensin II receptor 1 (AT1) blockade in cisplatin-induced nephrotoxicity in rats: Gender-related differences. *Ren Fail* 2012;34:1046-51.
25. Ghasemi M, Nematbakhsh M, Pezeshki Z, Soltani N, Moeini M, Talebi A. Nephroprotective effect of estrogen and progesterone combination on cisplatin-induced nephrotoxicity in ovariectomized female rats. *Indian J Nephrol* 2016;26:167-75.
26. Pezeshki Z, Nematbakhsh M, Mazaheri S, Eshraghi-Jazi F, Talebi A, Nasri H, *et al.* Estrogen abolishes protective effect of erythropoietin against cisplatin-induced nephrotoxicity in ovariectomized rats. *ISRN Oncol* 2012;2012:890310.
27. Nematbakhsh M, Pezeshki Z, Eshraghi-Jazi F, Ashrafi F, Nasri H, Talebi A, *et al.* Vitamin E, vitamin C, or losartan is not nephroprotectant against cisplatin-induced nephrotoxicity in presence of estrogen in ovariectomized rat model. *Int J Nephrol* 2012;2012:284896.
28. Mazaheri S, Nematbakhsh M, Bahadorani M, Pezeshki Z, Talebi A, Ghannadi AR, *et al.* Effects of fennel essential oil on cisplatin-induced nephrotoxicity in ovariectomized rats. *Toxicol Int* 2013;20:138-45.
29. Loh JM, Tran AL, Ji L, Groshen S, Daneshmand S, Schuckman A, *et al.* Baseline glomerular filtration rate and cisplatin-induced renal toxicity in urothelial cancer patients. *Clin Genitourinary Cancer* 2017. doi: 10.1016/j.clgc.2017.08.016.
30. Salek T, Vesely P, Bernatek J. Estimated glomerular filtration rate in oncology patients before cisplatin chemotherapy. *Klin Onkol* 2015;28:273-7.
31. Scott LA, Madan E, Valentovic MA. Influence of streptozotocin (STZ)-induced diabetes, dextrose diuresis and acetone on cisplatin nephrotoxicity in Fischer 344 (F344) rats. *Toxicology* 1990;60:109-25.
32. Fukushima K, Okada A, Oe H, Hirasaki M, Hamori M, Nishimura A, *et al.* Pharmacokinetic-pharmacodynamic analysis of cisplatin with hydration and mannitol diuresis: The contribution of urine cisplatin concentration to nephrotoxicity. *Eur J Drug Metab Pharmacokin* 2018;43:193-203.
33. McKibbin T, Cheng LL, Kim S, Steuer CE, Owonikoko TK, Khuri FR, *et al.* Mannitol to prevent cisplatin-induced nephrotoxicity in patients with squamous cell cancer of the head and neck (SCCHN) receiving concurrent therapy. *Support Care Cancer* 2016;24:1789-93.
34. Chen MF, Yang CM, Su CM, Hu ML. Vitamin C protects against cisplatin-induced nephrotoxicity and damage without reducing its effectiveness in C57BL/6 mice xenografted with Lewis lung carcinoma. *NutrCancer* 2014;66:1085-91.
35. Darwish MA, Abo-Youssef AM, Khalaf MM, Abo-Saif AA, Saleh IG, Abdelghany TM. Vitamin E mitigates cisplatin-induced nephrotoxicity due to reversal of oxidative/nitrosative stress, suppression of inflammation and reduction of total renal platinum accumulation. *J Biochem Mol Toxicol* 2017;31:1-9.
36. Durak I, Ozbek H, Karaayvaz M, Ozturk HS. Cisplatin induces acute renal failure by impairing antioxidant system in guinea pigs: Effects of antioxidant supplementation on the cisplatin nephrotoxicity. *Drug Chem Toxicol* 2002;25:1-8.
37. Konishi H, Fujiwara H, Itoh H, Shiozaki A, Arita T, Kosuga T, *et al.* Influence of magnesium and parathyroid hormone on cisplatin-induced nephrotoxicity in esophageal squamous cell carcinoma. *Oncol Lett* 2018;15:658-64.

38. Saito Y, Okamoto K, Kobayashi M, Narumi K, Furugen A, Yamada T, *et al.* Magnesium co-administration decreases cisplatin-induced nephrotoxicity in the multiple cisplatin administration. *Life Sci* 2017;189:18-22.
39. Saito Y, Okamoto K, Kobayashi M, Narumi K, Yamada T, Iseki K. Magnesium attenuates cisplatin-induced nephrotoxicity by regulating the expression of renal transporters. *Eur J Pharmacol* 2017;811:191-8.
40. Saito Y, Kobayashi M, Yamada T, Kasashi K, Honma R, Takeuchi S, *et al.* Premedication with intravenous magnesium has a protective effect against cisplatin-induced nephrotoxicity. *Support Care Cancer* 2017;25:481-7.
41. Kidera Y, Kawakami H, Sakiyama T, Okamoto K, Tanaka K, Takeda M, *et al.* Risk factors for cisplatin-induced nephrotoxicity and potential of magnesium supplementation for renal protection. *PloS One* 2014;9:e101902.
42. Yoshida T, Niho S, Toda M, Goto K, Yoh K, Umemura S, *et al.* Protective effect of magnesium preloading on cisplatin-induced nephrotoxicity: A retrospective study. *Japanese J Clin Oncol* 2014;44:346-54.
43. Soltani N, Nematbakhsh M, Eshraghi-Jazi F, Talebi A, Ashrafi F. Effect of oral administration of magnesium on Cisplatin-induced nephrotoxicity in normal and streptozocin-induced diabetic rats. *Nephrourol Mon* 2013;5:884-90.
44. Ashrafi F, Haghshenas S, Nematbakhsh M, Nasri H, Talebi A, Eshraghi-Jazi F, *et al.* The role of magnesium supplementation in cisplatin-induced nephrotoxicity in a rat model: No nephroprotectant effect. *Int J Prev Med* 2012;3:637-43.
45. Goren MP. Cisplatin nephrotoxicity affects magnesium and calcium metabolism. *Med Pediatr Oncol* 2003;41:186-9.