

An Updated Systematic Review and Meta-Analysis on Association of Serum Lipid Profile With Risk of Breast Cancer Incidence

Abstract:

Background: This meta-analysis was performed to investigate the effect of serum lipids on the risk of breast cancer incidence. **Methods:** PubMed, Web of Science, Scopus, and EMBASE were searched systematically from January 1998 to April 2019. Inclusion criteria were English observational studies (cohort or case-control) and the concentration of at least one of the lipid profile components (total cholesterol/triglycerides/low-density lipoprotein cholesterol/high-density lipoprotein cholesterol) measured before a diagnosis of breast cancer (BC). The studies were included in which the relative risk (RR) had been reported with 95% confidence intervals (CIs). A random-effects model was used. **Results:** A total of 25 studies were found, including 2,882,789 participants in cohort studies with 45,481 cases with BC, and 1983 BC cases and 2963 case-control studies. Combined RR of cohort studies for the highest versus lowest for the BC was LDL-C: 0.95 (95% CI: 0.89–1.01), triglycerides (TG): 0.95 (95% CI: 0.91–0.99; $P = 0.02$), total cholesterol (TC): 0.98 (95% CI: 0.91–1.05), and HDL-C: 0.86 (95% CI: 0.63–1.18). Combined RR of case-control studies for the highest versus lowest was LDL-C: 1.08 (95% CI: 0.78–1.48), TG: 1.73 (95% CI: 0.94–3.18), TC: 1.02 (95% CI: 0.80–1.29), and HDL-C: 0.79 (95% CI: 0.65–0.97). **Conclusions:** Based on the results, it can be concluded that only TG but not TC and/or LDL-C had a significant inverse association with the risk of BC incidence. HDL-C showed a significant protective effect against breast cancer in postmenopausal women and case-control studies.

Keywords: Meta-analysis, breast cancer, cholesterol, HDL, LDL, triglycerides

Introduction

The association of serum lipids (total cholesterol [TC], high-density lipoprotein cholesterol [HDL-C], low-density lipoprotein cholesterol [LDL-C], and triglycerides [TG]) with the risk and incidence of different cancers has been proposed by many researchers.^[1-4] Modifications in the serum cholesterol levels are related to the etiology of colorectal and breast carcinoma.^[5] It could be due to the important role of cholesterol as a precursor of steroid hormones.^[6] In a study that was conducted on mice with breast cancer (BC), it was revealed that the primary metabolites of cholesterol could increase the size of tumor and metastasis.^[7] A meta-analysis that was based only on cohort studies found an inverse relationship between TG levels in serum and the risk of BC, but TC and LDL did not show any significant relationship with BC risk and occurrence. They found the protective effect of

HDL among postmenopausal women.^[8] However, they did not consider the case-control studies^[9-11] and several large cohort studies^[12-16] that inevitably could introduce biases. Another meta-analysis reviewed the association of TC, HDL-C, and LDL-C levels with the BC risk and confirmed the inverse association between TC and HDL-C and the risk of BC,^[17] but did not investigate the effect of TG on BC. Given the different and controversial results obtained from various studies and meta-analyses about the effect of lipid profiles on the risk of BC, the effect of lipid profile on BC remains questionable. Therefore, the current meta-analysis was performed to systematically evaluate the association between individual lipid components (TC, HDL, LDL, and TG) and BC risk in a more comprehensive study taking into account both updated studies in this area after 2015 through April 2020 and also case-control studies that were not included in the previous meta-analysis by Haibo *et al.* 2015.^[8]

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Amerizadeh A, Vaseghi G, Farajzadegan Z, Asgary S. An up-dated systematic review and meta-analysis on association of serum lipid profile with risk of breast cancer incidence. *Int J Prev Med* 2022;13:142.

Atefeh Amerizadeh,
Golnaz Vaseghi¹,
Ziba Farajzadegan²,
Sedigheh Asgary

Isfahan Cardiovascular Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, ¹Applied Physiology Research center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, ²Department of Community Medicine, Isfahan University of Medical Sciences, Isfahan, Iran

Address for correspondence:
Prof. Sedigheh Asgary,
Department of Cardiovascular Research, Isfahan University of Medical Sciences, PO Box: 81465-1148, Isfahan, Iran.
E-mail: sedighehasgary@gmail.com

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.ir

DOI:
10.4103/ijpvm.IJPVM_285_20

Quick Response Code:



Material and Methods

These systematic review and meta-analysis were performed following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses statement recommendations.

Systematic search

Relevant studies were identified by searching PubMed, Scopus, Web of Science, and EMBASE from 1998 up to April 2019 using Medical Subject Heading [MeSH] and related keywords. We used AND and OR Boolean Operators to combine different concepts and similar concepts, respectively. The following keywords were used: “lipid,” “cholesterol,” “triglyceride,” “high-density lipoprotein [HDL],” “low-density lipoprotein [LDL],” “breast neoplasms,” “breast cancer,” “risk,” “incidence,” and “prevalence.” The search strategy in PubMed was “(lipid[Title/Abstract]) OR (cholesterol[Title/Abstract]) OR (HDL[Title/Abstract]) OR (LDL[Title/Abstract]) OR (triglycerides[Title/Abstract]) AND ((breast cancer[Title/Abstract]) OR (breast tumor[Title/Abstract])) AND (breast tumor[Title/Abstract]).”

We included prospective cohort and case-control studies estimating the relative risk (RR) with 95% confidence intervals (CIs) for the associations of specific lipid components including TC, HDL-C, LDL-C, and TG with breast cancer risk of incidence.

Study screening

Eligible studies for this meta-analysis were fulfilled in the following criteria: (1) the study design was an observational study (cohort or case-control study), (2) the exposure of interest was serum concentration of at least one of the selected lipid components (TC, HDL-C, LDL-C, and TG) measured before a diagnosis of BC, (3) the outcome of interest was the occurrence of BC, and (4) the relative risk (RR) with corresponding 95% CI or data to calculate it had been reported. If data had been duplicated in more than one study, we included the study with the most number of cases.

Quality assessment

Since all of the included studies were observational, their methodological quality was assessed by the nine-star Newcastle Ottawa Scale,^[18] which consists of three major aspects: selection, comparability, and exposure or outcome. A study with seven or more stars is considered to be high quality. Quality assessment was performed by two authors independently.

Data extraction

Two independent investigators (AA and GV) screened all retrieved articles and extracted the data from all eligible publications. Any disagreement was settled by discussion. In the end, a third researcher (ZF) approved the selection to ensure the accuracy of the data. The following data

were recorded: first author’s surname, publication year, country of origin, ethnicity, study and follow-up periods, characteristics of the study population (age and menopausal status), numbers of cases and participants, ranges of serum lipid levels, RRs from the most fully adjusted model for the highest versus lowest category of serum lipids, and the corresponding 95% CIs and matching or adjustments for confounding factors.

Statistical analysis

Two measures of association were used in the studies: Odds Ratio (OR) in the case-control and incidence rate ratio or Hazard Ratio (HR) in the cohort studies. For simplicity, we report the Relative Risk (RR) as a common measure of association. As the prevalence of BC is quite low, the OR in case-control and HR in cohort studies yielded a similar estimation of RR.^[19] In each study, the risk estimates reflected the comparison of the highest versus lowest categories. We produced some forest plots to evaluate the adjusted RRs and corresponding 95% CI visually across studies. If the heterogeneity among the studies was significant, the random-effects model using Der Simonian and Laird methods was used to summary RR and its corresponding 95% CI, otherwise, a model with fixed effect was calculated.^[20] The homogeneity among the studies was assessed with Cochran’s Q statistic and the I² statistic. For the I² statistic, values above 50%, and for the Chi-square test, *P* values < 0.1 were statistically supposed significant.^[21] We did subgroup analyses for each component to assess the impacts of different variables on outcomes and identify the sources of heterogeneity based on some characteristics including the geographic area, follow-up length, study design, and menopausal status. The sensitivity analysis was measured for assessment of the robustness of the combined risk estimates to evaluate whether the low-quality studies would influence the overall result. Publication bias was evaluated with Begg’s rank correlation tests and Egger’s linear regression tests.^[22,23] All analyses were performed using STATA version 14.

Results

Study characteristics

PubMed, Scopus, Web of Science, and EMBASE were searched for observational studies from 1998 to April 2019 and a total of 3544 articles were found. After removing duplicates, non-human studies, and non-English studies, 125 remained. After excluding the studies which did not meet our inclusion criteria finally, 25 studies were selected including 21 cohort studies^[13-16,24-40] and 4 case-control studies.^[9,11,12,41] The main characteristics of the selected studies have been summarized in Table 1. Studies were found and collected through the systematic search and selection process. Figure 1 shows the study selection procedure.

Altogether, 2,882,789 cases had participated in the cohort studies of which, 45,481 cases were affected with BC.

Table 1: The characteristics of the studies on serum lipids and breast cancer included in this meta-analysis

Author/Year	Country	Type of study	No. of cases/ participants	Age (years)	Follow up (years)	Outcome	Menopausal status	Study quality	Adjustments
Chandler <i>et al.</i> , 2016	United States	Cohort	864/15,602	>45	19	HDL, LDL, TC, TG	overall	9	Age, race, hormone replacement therapy, cigarette smoking, exercise, alcohol consumption, postmenopausal status, family history of cancer, aspirin use, history of fibrocystic or benign breast disease, total vegetable and fruit intake, history of mammogram, reproductive history including the age of menarche, oral contraceptive use, age, and BMI
Laamiri <i>et al.</i> , 2013	Moroccan	Case-control	Case: 400 Control: 400	45-83	2	HDL, LDL, TC, TG	overall	8	Age
Capasso <i>et al.</i> , 2010	Italy	Case-control	Case: 210 Control: 289	57.5	NA	HDL	Postmenopause	6	Tobacco use, alcohol abuse, food intake, physical activity grade, parity, age of menarche, menopausal status, oral contraceptive use, hormonal therapy use, personal and familial history of cancer
Furberg <i>et al.</i> , 2004	Norwegian	Cohort	708/38,823	58.4	17.2	HDL	Postmenopause, Premenopause	9	Age, BMI, county of residence, parity, height, serum TC, physical activity, blood pressure, serum TG, age at first birth, time since last meal, smoking, energy and fat intake, menopausal status
His <i>et al.</i> , 2014	France	Cohort	141/4433	49.5	11.5	HDL, LDL, TC, TG	overall	9	Age, BMI, intervention group, number of dietary records, alcohol intake, physical activity, smoking, education, height, family history of breast cancer, menopausal status, number of full-term, HRT use, energy intake, hyperlipidemia medication use, glycemia
His <i>et al.</i> , 2017	Switzerland	Cohort	583/1043	40-65	NA	HDL, LDL, TC, TG	overall	8	Age, BMI, menopausal status, history of breast cancer in first-degree relatives, personal history of benign breast disease, daily alcohol intake, daily glyceemic load, daily lipid intake daily energy intake without alcohol
Bjorge, <i>et al.</i> , 2010	England	Cohort	4862/287320	58	11	TC, TG	overall	8	Age, BMI, year of birth, smoking, glucose
Agnoli <i>et al.</i> , 2010	Italy	Case-control	Case: 236 Control: 556	35-69	13.5	HDL, TG	Postmenopause	9	Age, years from menopause, number of full-term pregnancies, age at first birth, oral contraceptives, hormone therapy, years of education, history of breast cancer in first degree relatives, breastfeeding, smoking, and alcohol consumption
Hoyer <i>et al.</i> , 1992	Danish	Cohort	51/5207	30-80	4-26	HDL, LDL, TC, TG	overall	8	Age, BMI, smoking, menopausal status, age at menarche, number of full-term, pregnancies, alcohol, and coffee consumption
Gaard <i>et al.</i> , 1994	Norwegian	Cohort	302/31,209	20-54	10.4	LDL	overall	9	Age, BMI, height, menopausal status, smoking

Contd...

Table 1: Contd...

Author/ Year	Country	Type of study	No. of cases/ participants	Age (years)	Follow up (years)	Outcome	Menopausal status	Study quality	Adjustments
Eliassan <i>et al.</i> , 2005	United States	Cohort	3177/71,921	42-69	<10	TC	overall	6	Age, BMI, age at menarche, parity, age at first birth, height, family history of breast cancer and BBD, alcohol consumption, physical activity, menopausal status, age at menopause, HRT use
Kucharaska <i>et al.</i> , 2008	United States	Cohort	359/7575	54-64	NA	HDL	Overall	9	Age, race, BMI, age at menarche, smoking, HRT use, age at menopause
Kabat <i>et al.</i> , 2009	United States	Cohort	165/4888	50-79	8	HDL, TG	Postmenopause	6	Age, education, race, BMI, oral contraceptive use, HRT use, age at menarche, age at first birth, age at menopause, alcohol, family history of breast cancer, history of breast biopsy, physical activity, energy intake, smoking, randomization of HRT, calcium plus vitamin D, and dietary modification trials, waist circumference, glucose, blood pressure
Inoue <i>et al.</i> , 2009	Japan	Cohort	120/18,176	40-69	10.2	HDL, TG	Overall	8	Age, study area, smoking, ethanol intake, serum TC
Iso <i>et al.</i> , 2009	Japan	Cohort	178/21,685	40-69	12.4	TC	Overall	9	Age, BMI, smoking, hypertension, diabetes, hyperlipidemia medication use, intake of total vegetable, coffee, and ethanol, public health center
Fagherazzi <i>et al.</i> , 2010	France	Cohort	2932/69,088	40-65	12	TC, TG	Overall, Premenopause	6	Age, intake of alcohol, total fat, and energy, oral contraceptives use, age at menarche, age at menopause, number of children, age at first pregnancy, family history of breast cancer, history of BBD, diabetes status, education, HRT use
Kitahara <i>et al.</i> , 2011	Korea	Cohort	3805/433,115	30-95	12.7	TC	Overall, Postmenopause, Premenopause	8	Age, BMI, smoking, alcohol intake, glucose, hypertension, physical activity
Bosco <i>et al.</i> , 2012	United States	Cohort	1228/49,172	21-69	10.5	TC	Overall, Postmenopause, Premenopause	7	Age, BMI, education, physical activity, obesity, Type 2 diabetes, hypertension
Melvin <i>et al.</i> , 2012	Swedish	Cohort	6105/234,494	>25	8.3	HDL, LDL, TC, TG	Overall	7	Age, parity, level of glucose, TG, TC, fasting status, and socioeconomic status
Strohmaier <i>et al.</i> , 2013	European	Cohort	5228/288,057	40.3-47.5	11.7	TC	Overall	8	Age, BMI, smoking
Ha <i>et al.</i> , 2009	Korea	Cohort	741/170,374	40- 64	8.84	TC	Postmenopause	8	Age, age at menarche, BMI, nulliparity, hormone replacement therapy, duration of breastfeeding, smoking habit, and alcohol consumption
Moorman <i>et al.</i> , 1998	United States	Case-control	Case: 196 Control:	NA	NA	HDL	Postmenopause, Premenopause	8	Education, smoking status, BMI, alcohol consumption, and history of benign breast disease

Contd...

Table 1: Contd...

Author/Year	Country	Type of study	No. of cases/ participants	Age (years)	Follow up (years)	Outcome	Menopausal status	Study quality	Adjustments
Osaki <i>et al.</i> , 2011	Japan	Cohort	78/15,386	55	9.1	HDL, TG	Overall	9	Age, smoking status, heavy drinking, presence of metabolic syndrome or pre-metabolic syndrome of each definition.
Borena <i>et al.</i> , 2010	European	Cohort	5006/256,512	44.2	11.9	TG	Overall	9	Baseline age, BMI, and smoking status
Manjer <i>et al.</i> , 2001	European	Cohort	409/9738	49.6	15	TC	Postmenopause, Premenopause	9	Age, nulliparity, current oral contraceptive use, current HRT, smoking, alcohol consumption, height, and weight

Moreover, 1983 cases and 2963 controls have participated in the case-control studies. Studies were published between 1998 and 2019, and their follow-up periods had a range from 2 to 13.5 years for the case-control and 4 to 26 years for the cohort studies. The populations were categorized into three groups: Americans (n = 6), Europeans (n = 13), Africans (n = 1), and Asians (n = 5). Except for two studies conducted among nurses and teachers,^[24,25] the rest were population-based. Of the 25 included studies, 15 contained the results for TC, 14 for HDL-C, 13 for TG, and 7 for LDL-C. Four studies only included postmenopausal women^[9,12,15,26] and 5 studies presented the estimates by menopausal status.^[24,27-29,40,41]

The results of the study quality assessment (score 0–9) ranged from 6 to 9, with an average score of 7.8, indicating high-quality studies.

Main results

HDL-C: Combined RR for the highest versus lowest HDL-C categories are shown in Figure 2. The summary result of 14 studies that examined the effect of HDL-C categories on BC was 0.86 (95% CI: 0.63–1.18). The negative association between HDL-C and BC in the case-control studies was significant at 0.79 (95% CI: 0.65–0.97), but the pooled risk estimate in the cohort studies was not 0.87 (95% CI: 0.58–1.32).

LDL-C: Seven studies examined the relationship between LDL-C and BC. As shown in Figure 3, the combined RR




Figure 1: Flow diagram of study selection. The flow chart shows literature search and selection for prospective cohort and case-control studies of serum lipids about the breast cancer risk. TC = total cholesterol; HDL-C = high-density lipoprotein cholesterol; LDL-C = low-density lipoprotein cholesterol; TG = triglycerides.




Figure 2: Forest plot of the highest versus lowest categories of serum HDL-C levels and breast cancer risk. Squares indicate study-specific relative risk estimates (size of the square reflects the study-specific statistical weight); horizontal lines indicate 95% confidence intervals (CI); diamond indicates the overall relative risk with its 95% confidence interval. a: Premenopausal; b: Postmenopausal

for the highest versus lowest LDL-C concentrations was 0.95 (95% CI: 0.89–1.01). The summary risk estimate of BC in five included cohort studies was 0.94 (95% CI: 0.88–1.01) and the combined result from two included case-control studies was 1.08 (95% CI: 0.78–1.48).

TG: An association between BC risk and serum TG was reported in 13 studies [Figure 4]. The pooled RR for the highest category versus the lowest was 1.15 (95% CI: 0.96–1.38). The result of meta-analysis on 10 cohort studies showed that the highest levels of TG compared to the lowest levels had a significant effect on the reduction of BC (RR = 0.95) (95% CI: 0.91–0.99; $P = 0.02$). Inversely, the combined results of the three case-control studies did not report a significant effect of 1.73 (95% CI: 0.94–3.18).

TC: In 15 studies, the effect of TC on BC had been examined. The summary risk estimate of BC for the highest TC compared with the lowest was 0.98 (95% CI: 0.91–1.05). The combination of the results of 15 cohort studies showed this effect size of 0.97 (95% CI: 0.90–1.06), and in the case-control studies ($n = 2$) pooled effect size of TC on BC was 1.02 (95% CI: 0.80–1.29). The result is shown in Figure 5.

Subgroup analysis

Subgroup analysis was performed according to some factors, such as the geographic region, follow-up length, the number of cases, and menopausal status. According to the results, when we stratified the analysis according to the geographical region, a significant positive effect of TC on BC in Asian studies (RR = 1.15, 95% CI: 1.03–1.28) and a positive relation between TG and




Figure 3: The forest plot of the highest vs. the lowest categories of serum LDL-C levels and breast cancer risk. Squares indicate study-specific relative risk estimates (size of the square reflects the study-specific statistical weight); The horizontal lines indicate 95% confidence intervals (CI); The diamond indicates the overall relative risk with its 95% confidence interval.

BC (RR = 3.78, 95% CI: 2.73–5.23) among African studies were detected [Table 2]. The stratified analysis according to menopausal status showed that an increase in HDL-C levels could cause a reduction in BC risk among women who were postmenopausal at baseline (RR = 0.84, 95% CI: 0.74–0.97), while a direct relationship between TG and BC risk in the postmenopausal women was found (RR = 1.43, 95% CI: 1.16–1.76). The result of the subgroup analysis based on case-control studies was significant only for TG. As a result, an inverse association between TG and BC risk was observed among the studies with more than 500 cases (RR = 0.94, 95% CI: 0.90–0.99). Inversely in studies with less than 500 cases, a direct relationship between TG and BC was detected (RR = 1.36, 95% CI: 1.01–1.84). We did not find any evidence of significant effects on BC in other subgroups.

Publication bias

Begg's rank correlation and Egger linear regression tests were used for the evaluation of publication bias for each lipid component. The obtained results for TC and LDL-C did not confirm the publication bias, while the outcomes of these tests were significant for HDL-C and TG (HDL-C: P value = 0.008, TG: P value = 0.015). Therefore, the Trim and fill methods were used to modify the results; however, the results did not change.

Sensitivity analysis

Sensitivity analysis was done to examine the effect of each study on the final result. Each time, one study was removed and the rest were separately analyzed for each lipid component. According to the HDL-C results, Hoyer *et al.* (1992)^[16] reported the most effective which caused a change in RR estimate of 0.09 (RR = 1.02, 95% CI: 0.92–




Figure 4: The forest plot of the highest vs. lowest categories of serum TG levels and breast cancer risk. Squares indicate study-specific relative risk estimates (size of the square reflects the study-specific statistical weight); The horizontal lines indicate 95% confidence intervals (CI); The diamond indicates the overall relative risk with its 95% confidence interval. a: Premenopausal; b: Postmenopausal

1.15). Laamiri *et al.* (2013)^[11] also caused a change in RR estimate of 0.09 on TG which with the elimination of this study, the result changed but not significantly (RR = 1.02, 95% CI: 0.88–1.17). The sensitivity analysis on TC showed that three studies, Bosco *et al.* (2012), Bjørge *et al.* (2010), and Fagherazzi *et al.* (2010),^[25,29,38] caused changes in RR estimate of 0.02. So we removed these studies and analyzed the remaining studies. But the result was not still significant (RR = 1, 95% CI: 0.92–1.08). For LDL-C, it was found that studies did not affect the combined risk estimates.

Discussion

In the current meta-analysis, 21 cohort studies^[13-16,24-39] and 4 case-control studies^[9,11,12,40] were pooled and analyzed. Altogether, 45,481 BC cases in cohort studies and 1983 cases in the case-control studies were studied. So far, the most up-to-date and comprehensive meta-analysis on the association of lipid profile with breast cancer risk was published by Haibo *et al.* 2015,^[8] which included 15 cohort studies. In our meta-analysis on this topic, 21 cohort studies and 4 case-control studies^[9,11,12,40] were gathered and analyzed in addition to their references. More than two articles that were published between April 2015 and April 2019^[31,32] were also reviewed in our meta-analysis. They excluded Borena *et al.* 2011^[14] and Ha *et al.* 2009^[15] because they believed that they duplicated the reports of Strohmaier *et al.*, 2013,^[34] and Kitahara *et al.*, 2011,^[33] respectively. When we went, however, through the articles more deeply, we found that although the population studied by Borena *et al.* 2011^[14] and Strohmaier *et al.* 2013^[34]




Figure 5: The forest plot of the highest vs. lowest categories of serum TC levels and breast cancer risk. Squares indicate study-specific relative risk estimates (size of the square reflects the study-specific statistical weight); The horizontal lines indicate 95% confidence intervals (CI); The diamond indicates the overall relative risk with its 95% confidence interval. a: Premenopausal; b: Postmenopausal

was the same (Norway, Austria, and Sweden), the former had studied the role of TG in cancer risk but the later studied the role of cholesterol on cancer risk. Although the population studied by Ha *et al.* 2009^[15] is a subset of a larger one by Kitahara *et al.* 2011,^[33] we considered both in our study because we wanted to study the postmenopausal women separately in our subgroup analysis.

According to the results, all lipid profile components were related to the risk of BC occurrence; however, the association was not significant. The subgroup analysis showed a significant association only for TG. Moreover, an inverse association between TG and BC risk was observed among the studies with more than 500 cases. A significant positive effect of TC was also observed among the studies in Asia and a positive relation between TG and BC in a study in Africa.

According to the subgroup analysis, HDL-C showed a positive protective effect among postmenopausal women. A negative association between HDL-C and BC was also found in the case-control studies. These results are in agreement with reports by Esposito *et al.* 2013^[41] and Haibo *et al.* 2015^[8] about the protective role of HDL-C against BC. The protective effect of HDL-C on BC occurrence can be explained by the role of serum HDL-C level as a marker of androgen status.^[41] Among postmenopausal women, the ovaries and adrenals keep on producing androgens which can be converted to biologically active estrogens in peripheral tissues. Therefore, the lack of HDL-C might result in androgen deficiency and subsequent

Table 2: The stratified meta-analyses of lipid components and breast cancer risk

Group	TC				HDL-C			
	No. of studies	RR (95%CI)	I ² (%)	P*	No. of studies	RR (95%CI)	I ² (%)	P*
Total	20	0.98 (0.91-1.05)	64.5	0.504	17	0.93 (0.69-1.25)	95.5	0.611
Study design								
Cohort	18	0.97 (0.90-1.05)	68	0.476	10	0.78 (0.58-1.32)	97.3	0.521
Case-control	2	1.02 (0.80-1.29)	0	0.977	7	1.11 (0.93-1.34)	34.4	0.254
Menopausal status								
Premenopausal	4	1 (0.85-1.18)	0	0.982	4	0.95 (0.57-1.59)	67	0.853
Postmenopausal	4	1.09 (0.99-1.20)	0	0.083	7	1.01 (0.77-1.31)	74.1	0.959
Geographic area								
Unaited states	6	1 (0.95-1.05)	0	0.999	8	1 (0.85-1.18)	43.2	0.273
Europe	10	0.89 (0.78-1.02)	69.9	0.090	10	0.82 (0.48-1.39)	97.3	0.208
Asia	3	1.15 (1.03-1.28)	0	0.013	2	1.41 (1.01-1.97)	0	0.045
Africa	1	1.12 (0.72-1.74)	-	0.613	1	0.85 (0.28-2.58)	-	0.510
Length of follow-up								
≥10 year	14	0.96 (0.86-1.06)	73.9	0.199	10	0.78 (0.50-1.21)	97.4	0.271
<10 year	5	1 (0.94-1.07)	0	0.944	4	1.02 (0.86-1.21)	0	0.814
No. of cases								
≥500	14	0.97 (0.90-1.05)	72.7	0.433	5	0.96 (0.89-1.02)	52.9	0.191
<500	6	1.04 (0.84-1.30)	7.4	0.711	12	0.91 (0.57-1.47)	96.4	0.710
Group	LDL-C				TG			
	No. of studies	RR (95%CI)	I ² (%)	P*	No. of studies	RR (95%CI)	I ² (%)	P*
Total	7	0.95 (0.89-1.01)	0	0.107	16	1.11 (0.94-1.30)	92.4	0.226
Study design								
Cohort	5	0.94 (0.88-1.01)	0	0.077	12	0.95 (0.91-0.99)	0	0.019
Case-control	2	1.08 (0.78-1.48)	17.5	0.654	4	1.45 (0.89-2.36)	96.4	0.137
Menopausal status								
Premenopausal	-	-	-	-	4	1.17 (0.80-1.69)	84.9	0.415
Postmenopausal	-	-	-	-	2	0.84 (0.66-1.06)	0	0.136
Geographic area								
Unaited states	1	0.95 (0.88-1.02)	-	0.169	5	0.92 (0.82-1.04)	42	0.170
Europe	5	0.90 (0.78-1.05)	0	0.185	10	1.04 (0.85-1.28)	92.9	0.708
Asia	-	-	-	-	2	0.92 (0.63-1.34)	0	0.660
Africa	1	1.26 (0.84-1.88)	-	0.261	1	3.78 (2.73-5.23)	-	<0.001
Length of follow-up								
≥10 year	4	0.94 (0.84-1.04)	6	0.229	13	0.99 (0.84-1.18)	90.7	0.950
<10 year	2	1.02 (0.76-1.37)	42.2	0.872	4	1.38 (0.66-2.89)	95.8	0.392
No. of cases								
≥500	3	0.95 (0.88-1.01)	0	0.106	6	0.94 (0.90-0.98)	0	0.006
<500	4	0.98 (0.78-1.22)	40.7	0.827	10	1.28 (0.95-1.73)	89.3	0.098

* statistically significant in the level of 0.05. TC: total cholesterol TG: triglycerides. LDL-C: Low-density lipoprotein cholesterol. HDL-C: High-density lipoprotein cholesterol

BC development. It is also believed that HDL-C has anti-inflammatory properties which are inversely associated with BC risk.^[42]

Dyslipidaemia is considered responsible for BC.^[42,43] The carcinogenic potential of TG in tumor growth as an independent source for fatty acid oxidation has been stated previously.^[44,45] Recently, it was found that body mass index (BMI) and obesity increase the chance of triple-negative breast cancer (TNBC) occurrence.^[45-47] An association has also been reported between the increased risk of BC recurrence and mortality with obesity and high BMI.^[46,47]

The results on the effect of TG on BC were also conflicting. TG was found to have a positive effect on BC among the studies with less than 500 cases. Meanwhile, despite triglyceride carcinogenic potential, a significant inverse relationship was observed between TG and BC risk in the cohort studies, but not among case-control studies with more than 500 cases. In line with our findings, Haibo *et al.* 2015^[8] also proposed that TG may be inversely associated with BC risk. The biological mechanisms for the inverse association between TG and BC risk still remain unclear. Meanwhile, in our analysis by ethnicity, we observed a significant positive relationship between TG and BC risk in African but not in Asian, European, and American populations. Since only one study for Africa was available, more studies are necessary to determine whether this association is valid.

A positive effect of cholesterol on BC risk was observed among Asian populations in five studies included in this meta-analysis. These findings are in agreement with the ethnicity analysis by Haibo *et al.* 2015,^[8] because they also reported a positive effect of cholesterol on BC risk in two Asian studies. Accordingly, it seems that the ethnicity difference might have an important role in heterogeneity. The positive role of cholesterol in BC progression has been attributed to its role as a precursor of steroid hormones, estrogen, and progesterone.^[48,49] Cholesterol has been also reported to accelerate and enhance tumor growth and formation. Some epidemiological studies showed that there is an association between abnormal levels of HDL-C and LDL-C in patients with cancer. However, no positive correlation was observed in our ethnicity analysis between cholesterol levels and the risk of BC except for the Asian population. These discrepancies among different ethnicities could be attributed to genetic variations and high heterogeneity of the BC subtypes. The metabolic heterogeneity which is likely to be present in all cancers, including BC, can be also considered as an explanation for this association.^[49,50]

LDL-C level showed no significant effect on BC in our analysis. Meanwhile, LDL-C lowering drugs such as statins have been widely studied for their positive role on BC.^[51] Inversely, some studies presented no association between

statin use and the BC risk.^[52] Mansourian *et al.* (2016)^[53] merged the results for 124,669 cases (eight cohort studies) with BC. Their results suggest a significant reduction in recurrence and death among statin users. On the other hand, Undela *et al.* (2012)^[54] studied 24 studies (76,759 participants), and finally, they did not find any significant reduction in BC incidence among statin users, either short- or long-term. In this study, LDL also showed no significant effects on BC risk even in subgroup analysis. Since statins work through LDL reduction, our results are supported by Undela *et al.* (2012)'s^[54] study. LDL was the only lipid profile content that did not show a significant heterogeneity among the studies. TC was the second with less heterogeneity but TG and HDL showed substantial heterogeneity among the studies. For HDL-C, Hoyer *et al.* (1992)^[16] presented the most effective, and for TG, Laamiri *et al.* (2013)^[11] showed significant heterogeneity. Future research with larger sample sizes, higher quality, detailed information on geographical regions and menopausal status, and less variation in methodology is warranted to extend our findings.

Conclusions

According to this meta-analysis, an inverse association was observed between breast carcinogenesis risk and the serum TG in cohort-design and case-control studies with a population of more than 500. Serum HDL-C showed almost a protective role against breast cancer among postmenopausal women and case-control studies.

Financial support and sponsorship

This study was supported financially by Isfahan University of Medical Sciences.

Conflicts of interest

There are no conflicts of interest.

Received: 19 May 20 **Accepted:** 13 Sep 21

Published: 23 Nov 22

References

1. Schatzkin A, Hoover RN, Taylor PR, Ziegler RG, Carter CL, Albanes D, *et al.* Site-specific analysis of total serum cholesterol and incident cancer in the national health and nutrition examination survey I epidemiologic follow-up study. *Cancer Res* 1988;48:452-8.
2. Gerber M, Richardson S, Crastes de Paulet P, Pujol H, Crastes de Paulet A. Relationship between vitamin E and polyunsaturated fatty acids in breast cancer. Nutritional and metabolic aspects. *Cancer* 1989;64:2347-53.
3. Chyou PH, Nomura AM, Stemmermann GN, Kato I. Prospective study of serum cholesterol and site-specific cancers. *J Clin Epidemiol* 1992;45:287-92.
4. Forones NM, Falcao JB, Mattos D, Barone B. Cholesterolemia in colorectal cancer. *Hepatogastroenterology* 1998;45:1531-4.
5. Patel PS, Shah MH, Jha FP, Raval GN, Rawal RM, Patel MM, *et al.* Alterations in plasma lipid profile patterns in head and neck cancer and oral precancerous conditions. *Indian J Cancer* 2004;41:25-31.

6. Li X, Tang H, Wang J, Xie X, Liu P, Kong Y, *et al.* The effect of preoperative serum triglycerides and high-density lipoprotein-cholesterol levels on the prognosis of breast cancer. *The Breast* 2017;32:1-6.
7. Nelson ER, Wardell SE, Jasper JS, Park S, Suchindran S, Howe MK, *et al.* Hydroxycholesterol links hypercholesterolemia and breast cancer pathophysiology. *Science* 2013;342:1094-8.
8. Haibo N, Huixiang L, Rong G. Serum lipids and breast cancer risk: A meta-analysis of prospective cohort studies. *PLoS One* 2015;10:e0142669.
9. Agnoli C, Berrino F, Abagnato CA, Muti P, Panico S, Crosignani P, *et al.* Metabolic syndrome and postmenopausal breast cancer in the ORDET cohort: A nested case-control study. *Nutr Metab Cardiovasc Dis* 2010;20:41-8.
10. Martin LJ, Melnichouk O, Huszti E, Connelly PW, Greenberg CV, Minkin S, *et al.* Serum lipids, lipoproteins, and risk of breast cancer. A nested case-control study using multiple time points. *J Natl Cancer Inst* 2015;107:djv032.
11. Laamiri FZ, Otmani A, Ahid S, Barkat A. Lipid profile among Moroccan overweight women and breast cancer: A case-control study. *Int. J. Gen. Med.* 2013;6:439-45.
12. Capasso I, Esposito E, Pentimalli F, Crispo A, Montella M, Grimaldi M, *et al.* Metabolic syndrome affects breast cancer risk in postmenopausal women: National Cancer Institute of Naples experience. *Cancer Biol Ther* 2010;10:1240-3.
13. Osaki Y, Taniguchi S, Tahara A, Okamoto M, Kishimoto T. Metabolic syndrome and incidence of liver and breast cancers in Japan. *Cancer Epidemiol* 2012;36:141-7.
14. Borena W, Stocks T, Jonsson H, Strohmaier S, Nagel G, Bjørge T, *et al.* Serum triglycerides and cancer risk in the metabolic syndrome and cancer (Me-Can) collaborative study. *Cancer Causes Control* 2011;22:291-9.
15. Ha M, Sung J, Song YM. Serum total cholesterol and the risk of breast cancer in postmenopausal Korean women. *Cancer Causes Control* 2009;20:1055-60.
16. Høyer AP, Engholm G. Serum lipids and breast cancer risk: A cohort study of 5,207 Danish women. *Cancer Causes Control* 1992;3:403-8.
17. Touvier M, Fassier P, His M, Norat T, Chan DS, Blacher J, *et al.* Cholesterol and breast cancer risk: A systematic review and meta-analysis of prospective studies. *Br J Nutr* 2015;114:347-57.
18. Stang A. Critical evaluation of the Newcastle-Ottawa scale for the assessment of the quality of nonrandomized studies in meta-analyses. *Eur J Epidemiol* 2010;25:603-5.
19. Greenland S, Longnecker MP. Methods for trend estimation from summarized dose-response data, with applications to meta-analysis. *Am J Epidemiol* 1992;135:1301-9.
20. Lau J, Ioannidis JP, Schmid CH. Quantitative synthesis in systematic reviews. *Ann Intern Med* 1997;127:820-6.
21. Higgins JP, Thompson SG, Deeks JJ, Altman DG. Measuring inconsistency in meta-analyses *BMJ* 2003;327:557-60.
22. Egger M, Davey Smith G, Schneider M, Minder C. Bias in meta-analysis detected by a simple, graphical test. *BMJ* 1997;315:629-34.
23. Begg CB, Mazumdar M. Operating characteristics of a rank correlation test for publication bias. *Biometrics* 1994;50:1088-101.
24. Eliassen AH, Colditz GA, Rosner B, Willett WC, Hankinson SE. Serum lipids, lipid-lowering drugs, and the risk of breast cancer. *Arch Intern Med* 2005;24:165:2264-71.
25. Fagherazzi G, Fabre A, Boutron-Ruault MC, Clavel-Chapelon F. Serum cholesterol level, use of a cholesterol-lowering drug, and breast cancer: Results from the prospective E3N cohort. *Eur J Cancer Prev* 2010;19:120-5
26. Kabat GC, Kim M, Chlebowski RT, Khandekar J, Ko MG, McTiernan A, *et al.* A longitudinal study of the metabolic syndrome and risk of postmenopausal breast cancer. *Cancer Epidemiol Biomarkers Prev* 2009;18:2046-53.
27. Furberg AS, Veierød MB, Wilsgaard T, Bernstein L, Thune I. Serum high-density lipoprotein cholesterol, metabolic profile, and breast cancer risk. *J Natl Cancer Inst* 2004;96:1152-60.
28. His M, Zelek L, Deschasaux M, Pouchieu C, Kesse-Guyot E, Hercberg S, *et al.* Prospective associations between serum biomarkers of lipid metabolism and overall, breast and prostate cancer risk. *Eur J Epidemiol* 2014;29:119-32.
29. Bosco JL, Palmer JR, Boggs DA, Hatch EE, Rosenberg L. Cardiometabolic factors and breast cancer risk in U.S. black women. *Breast Cancer Res Treat* 2012;134:1247-56.
30. Manjer J, Kaaks R, Riboli E, Berglund G. Risk of breast cancer in relation to anthropometry, blood pressure, blood lipids and glucose metabolism: A prospective study within the Malmö Preventive Project. *Eur J Cancer Prev* 2001;10:33-42.
31. Chandler PD, Song Y, Lin J, Zhang S, Sesso HD, Mora S, *et al.* Lipid biomarkers and long-term risk of cancer in the Women's Health Study. *Am J Clin Nutr* 2016;103:1397-407.
32. His M, Dartois L, Fagherazzi G, Boutten A, Dupré T, Mesrine S, *et al.* Associations between serum lipids and breast cancer incidence and survival in the E3N prospective cohort study. *Cancer Causes Control* 2017;28:77-88.
33. Kitahara CM, Berrington de González A, Freedman ND, Huxley R, Mok Y, Jee SH, *et al.* Total cholesterol and cancer risk in a large prospective study in Korea. *J Clin Oncol* 2011;29:1592-8.
34. Strohmaier S, Edlinger M, Manjer J, Stocks T, Bjørge T, Borena W, *et al.* Total serum cholesterol and cancer incidence in the Metabolic syndrome and Cancer Project (Me-Can). *PLoS One* 2013;8:e54242.
35. Inoue M, Noda M, Kurahashi N, Iwasaki M, Sasazuki S, Iso H, *et al.* Impact of metabolic factors on subsequent cancer risk: Results from a large-scale population-based cohort study in Japan. *Eur J Cancer Prev* 2009;18:240-7.
36. Iso H, Ikeda A, Inoue M, Sato S, Tsugane S, JPHC Study Group. Serum cholesterol levels in relation to the incidence of cancer: The JPHC study cohorts. *Int J Cancer* 2009;125:2679-86.
37. Kucharska-Newton AM, Rosamond WD, Mink PJ, Alberg AJ, Shahar E, Folsom AR. HDL-cholesterol and incidence of breast cancer in the ARIC cohort study. *Ann Epidemiol* 2008;18:671-7.
38. Bjørge T, Lukanova A, Jonsson H, Tretli S, Ulmer H, Manjer J, *et al.* Metabolic syndrome and breast cancer in the Me-Can (Metabolic Syndrome and Cancer) project. *Cancer Epidemiol Biomarkers Prev* 2010;19:1737-45.
39. Melvin JC, Seth D, Holmberg L, Garmo H, Hammar N, Jungner I, *et al.* Lipid profiles and risk of breast and ovarian cancer in the Swedish AMORIS study. *Cancer Epidemiol Biomarkers Prev* 2012;21:1381-4.
40. Moorman PG, Hulka BS, Hiatt RA, Krieger N, Newman B, Vogelman JH, *et al.* Association between high-density lipoprotein cholesterol and breast cancer varies by menopausal status. *Cancer Epidemiol Biomarkers Prev* 1998;7:483-8.
41. Esposito K, Chiodini P, Capuano A, Bellastella G, Maiorino MI, Rafaniello C, *et al.* Metabolic syndrome and postmenopausal breast cancer: Systematic review and meta-analysis. *Menopause* 2013;20:1301-9.
42. Sarrel PM. Cardiovascular aspects of androgens in women. *Semin Reprod Endocrinol* 1998;16:121-8.
43. Cibeira GH, Giacomazzi J, Aguiar E, Schneider S, Etrich B, DE Souza CI, *et al.* Apolipoprotein E genetic polymorphism,

- serum lipoprotein levels and breast cancer risk: A case-control study *Mol Clin Oncol* 2014;2:1009-15.
44. Saavedra-García P, Nichols K, Mahmud Z, Fan LY, Lam EW. Unravelling the role of fatty acid metabolism in cancer through the FOXO3-FOXO1 axis. *Mol Cell Endocrinol* 2017;462:82-92.
 45. Agresti R, Meneghini E, Baili P, Minicozzi P, Turco A, Cavallo I, *et al.* Association of adiposity, dysmetabolisms, and inflammation with aggressive breast cancer subtypes: A cross-sectional study. *Breast Cancer Res Treat* 2016;157:179-89.
 46. Gershuni V, Li YR, Williams AD, So A, Steel L, Carrigan E, *et al.* Breast cancer subtype distribution is different in normal weight, overweight, and obese women. *Breast Cancer Res Treat* 2017;163:375-81.
 47. Duggan C, Irwin ML, Xiao L, Henderson KD, Smith AW, Baumgartner RN, *et al.* Associations of insulin resistance and adiponectin with mortality in women with breast cancer. *J Clin Oncol* 2011;29:32-9.
 48. Warner M, Gustafsson JA. On estrogen, cholesterol metabolism, and breast cancer. *N Engl J Med* 2014;370:572-3.
 49. Kim Y, Park SK, Han W, Kim DH, Hong YC, Ha EH, *et al.* Serum high-density lipoprotein cholesterol and breast cancer risk by menopausal status, body mass index, and hormonal receptor in Korea. *Cancer Epidemiol Biomarkers Prev* 2009;18:508-15.
 50. Martinez-Outschoorn UE, Peiris-Page M, Pestell RG, Sotgia F, Lisanti MP. Cancer metabolism: A therapeutic perspective. *Nat Rev Clin Oncol* 2017;14:11-31.
 51. Cauley JA, McTiernan A, Rodabough RJ, LaCroix A, Bauer DC, Margolis KL, *et al.* Statin use and breast cancer: Prospective results from the Women's Health Initiative. *J Natl Cancer Inst* 2006;98:700-7.
 52. Boudreau DM, Yu O, Miglioretti DL, Buist DS, Heckbert SR, Daling JR. Statin use and breast cancer risk in a large population-based setting. *Cancer Epidemiol Biomarkers Prev* 2007;16:416-1.
 53. Mansourian M, Haghjooy-Javanmard S, Eshraghi A, Vaseghi G, Hayatshahi A, Thomas J. Statins use and risk of breast cancer recurrence and death: A systematic review and meta-analysis of observational studies. *J Pharm Sci* 2016;19:72-81.
 54. Undela K, Srikanth V, Bansal D. Statin use and risk of breast cancer: A meta-analysis of observational studies. *Breast Cancer Res Treat* 2012;135:261-9.