

The Prevalence of Cancer in Patients with Multiple Sclerosis (MS) who were Under Treatment with Natalizumab (Tysabri): A Systematic Review and Meta-Analysis

Abstract

Background: To determine the pooled prevalence of cancer in subjects with multiple sclerosis (MS) who received Natalizumab. **Methods:** Two researchers systematically searched PubMed, Scopus, EMBASE, Web of Science, google scholar, and gray literature including references of the included studies. The search strategy which was used in PubMed was (“Disseminated Sclerosis” OR “multiple sclerosis” OR “MS” OR “Acute Fulminating”) AND (“Cancer” OR “Neoplasia*” OR “Neoplasm*” OR “Tumor*” OR “Malignancy” OR “Benign Neoplasm” OR “Malignant neoplasm”) AND (“Tysabri” OR “Antegren” OR “natalizumab” OR “Modifying Therapy”). **Results:** We found 1,993 articles by literature search, and 1,573 studies remained after removing duplicate studies. For metaanalysis, we used the extracted data of eight studies. The pooled prevalence of cancer in patients who received Natalizumab was 2% (95%CI: 1–3%; I^2 : 99.4%, $P < 0.001$). The pooled prevalence of basal cell carcinoma in patients with cancer was 12% (95%CI: 5–20%; I^2 : 50.3%, $P = 0.13$). **Conclusions:** The main finding of this systematic review and metaanalysis is that the pooled prevalence of cancer in subjects who suffer from MS and received natalizumab was 2%.

Keywords: Multiple sclerosis, neoplasm, prevalence

Introduction

Multiple sclerosis (MS) is an inflammatory, autoimmune disease of central nervous system (CNS) that affects youth all over the world.^[1] The most common form of the disease is relapsing-remitting (RR) that accounts for nearly 85% of all MS types.^[2] Interferon-beta and glatiramer acetate are the first disease-modifying therapies for patients with MS while their effectiveness is not high.^[3] Natalizumab is a monoclonal antibody, an $\alpha 4$ -integrin antagonist, is a disease-modifying therapy (DMT) which is administered in subjects with MS.^[4] Natalizumab prevents migration of lymphocytes across the blood–brain barrier.^[4] It is widely used for treating RR form of MS disease.^[5,6] Its safety and efficacy profile is acceptable while concern regarding developing progressive multifocal leukoencephalopathy (PML) raises during treatment with this medication.^[7]

Subjects with MS have higher rate of mortality compared with their age- and sex-matched controls due to secondary complications of MS.^[8,9] Infectious,

respiratory, and cardiovascular diseases as well as malignancies are the most frequent leading causes of death in individuals with MS.^[10,11]

Smoking, vitamin D deficiency, and genetics are among common etiological factors between cancers and MS.^[12]

Previous studies reported different rates of cancer in MS patients who are under treatment with various medications such as Natalizumab.

So, we designed this systematic review and meta-analysis to estimate the pooled prevalence of cancer in MS patients who received Natalizumab.

Methods

The protocol of this systematic review was approved in Tehran University of medical sciences (IR.TUMS.NI.REC.1400.053).

The search terms were:

We systematically and comprehensively searched PubMed, Scopus, EMBASE, Web of Science, google scholar, and gray

How to cite this article: Nasirzadeh A, Jahanshahi R, Ghajarzadeh M, Mohammadi A, Sahraian MA, Moghadasi AN. The prevalence of cancer in patients with multiple sclerosis (MS) who were under treatment with natalizumab (Tysabri): A systematic review and meta-analysis. *Int J Prev Med* 2023;14:114.

Amirreza
Nasirzadeh,
Reza Jahanshahi¹,
Mahsa
Ghajarzadeh^{2,3},
Aida Mohammadi³,
Mohammad
Ali Sahraian²,
Abdorreza Naser
Moghadasi²

Student Research Committee, Mashhad University of Medical Sciences, Mashhad, Iran, ¹Golestan University of Medical Sciences, Gorgan, Iran, ²Multiple Sclerosis Research Center, Neuroscience Institute, Tehran University of Medical Sciences, Tehran, Iran, ³Universal Council of Epidemiology (UCE), Universal Scientific Education and Research Network (USERN), Tehran University of Medical Sciences, Tehran, Iran

Address for correspondence:

Dr. Abdorreza Naser Moghadasi, Sina Hospital, Tehran, Iran. E-mail: abdorrezamoghadasi@gmail.com

Access this article online

Website: www.ijpvmjournal.net/www.ijpvm.net

DOI: 10.4103/ijpvm.ijpvm_77_22

Quick Response Code:



This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

literature including references of the included studies that were published before October 2021.

("Disseminated Sclerosis" OR "multiple sclerosis" OR "MS" OR "Acute Fulminating") AND ("Cancer" OR "Neoplasia*" OR "Neoplasm*" OR "Tumor*" OR "Malignancy" OR "Benign Neoplasm" OR "Malignant neoplasm") AND ("Tysabri" OR "Antegren" OR "natalizumab" OR "Modifying Therapy").

We included studies if their design was cohort and providing information regarding the number of included patients who were treated with Natalizumab and incident number of cancer and also studies which were published in English language.

We excluded studies which were case-control, cross-sectional studies, and case reports.

Data extraction

Two researchers extracted data regarding the total number of patients, first author name, publication year, country of origin, and number of patients with cancer from the included studies.

If discrepancies were found, the third one reviewed the forms.

Risk of bias assessment

We assessed the potential risk of bias in included studies was evaluated using the Hoy assessment scale.^[13]

Statistical analysis

We conducted statistical analyses were performed using STATA (Version 13.0; Stata Corp LP, College Station, TX, USA). We used the inverse variance with random effects model. To determine heterogeneity, Inconsistency (I²) was calculated.

Results

We found 1,993 articles by literature search, and after deleting duplicates 1,573 remained. Eight articles remained for metaanalysis [Figure 1].

Eight articles were included. The basic characteristics of the included studies [Table 1].

The pooled prevalence of cancer in patients who received Natalizumab was 2% (95%CI: 1–3%; I²:99.4%, P < 0.001) [Figure 2].

The pooled prevalence of basal cell carcinoma in patients with cancer was 12% (95%CI: 5–20%; I²:50.3%, P = 0.13) [Figure 3].

The quality assessment of included studies are reported in Table 2.

Discussion

To the best of our knowledge, this is the first study




Figure 1: Flow diagram summarizing the selection of eligible studies




Figure 2: The pooled prevalence of basal cell carcinoma in cases who received Natalizumab

Evaluating the pooled prevalence of cancer in MS cases who received natalizumab. The results show that the pooled prevalence is 2% ranging from 0 to 5%. This finding could show that cancers should be evaluated in subjects with MS who receive Natalizumab. As it was mentioned in the introduction, MS is not a fatal disease and most cases die due to complications such as infections, respiratory diseases, and cancers.^[10,11]

Recent innovation by disease modifying therapies such as Natalizumab has led to better clinical progress, and quality of life improvement.^[14] Better anti-inflammatory responses and neuroprotective effects are advantages of Natalizumab treatment,^[14] but hepatotoxicity, allergic reactions, progressive multifocal leukoencephalopathy (PML) development, and a higher risk of infection are among disadvantages of Natalizumab treatment.^[15]

Previous studies showed risk of developing cancer in subjects who received Natalizumab is not higher.^[4,16]

Basal cell carcinoma was reported by most studies and its pooled prevalence was 12%.

In general population, breast, colorectal, prostate, lung, and stomach cancers are the most frequent neoplasm, and breast, cervical, and thyroid are the most common cancers in women.^[17] In MS cases, the most incident cancers reported as breast, and digestive cancers.^[12]

Foley *et al.*^[18] enrolled 6,634 MS patients who received natalizumab and reported malignant incidence as 449.0 per 100,000 patient-years. The most prevalent cancers among women and men in their study were breast and colon cancers, respectively.

In another large safety study, Stamatellos *et al.*^[19] enrolled 56,767 MS patients who were under treatment with natalizumab and reported cancer in 2.3%.

In another study, Polman *et al.*^[20] included 627 MS cases who were treated by natalizumab and found malignancies in 5.




Figure 3: The pooled prevalence of basal cell carcinoma in patients with cancer

Natalizumab is effective in patients with RR type of the disease while has multifocal leukoencephalopathy (PML) as its complication.^[6] PML is life threatening, so monitoring of JCV antibody status is necessary for patients who administer this medication.

Natalizumab is a monoclonal antibody which is approved for treating RR form of MS which slows down the progression of symptoms and decreases the rate of flare up.^[21] It decreases inflammation by blocking the cross of leukocytes from the blood vessel which leads to inflammation decrease.^[21]

It is shown that natalizumab plays role in rapid progression of the CNS diffuse large B-cell lymphomas (CNSL)^[16] and also some modifications in pigmented lesions.^[22] The incidence of melanoma in cases who received natalizumab estimated as 5/100,000 MS person-years.^[22]

Overall, it seems that administration of natalizumab is not related with higher risk of cancer in subjects with MS.

This systematic review has some limitation. First, all studies did not report the prevalence of each neoplasm separately. Second, the number of included studies was limited. Third, there was no exact data regarding the time between the availability and affordability of the drug and the incidence of cancers.

Conclusions

The main finding of this systematic review and metaanalysis is that the pooled prevalence of cancer in subjects who suffer from MS and received natalizumab was 2%.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

Received: 06 Mar 22

Accepted: 27 Oct 22

Published: 28 Sep 23

Table 1: Basic characteristics of included studies

Table X. Characteristics of included studies.																	
Author	Year	Country	Female	Male	Total sample	MS	Use natalizumab	NTB cancer	BCC cancer	LTC	OCPC	SCC	UN/NM	FBC	CC	MM	CIN
Alping	2020	Sweden	5287	849	6136		1670	44	8				17	2			15
DAmico	2019	Italy	792	388	1180		142	0									
Foley	2019	Multi-country	4749	1759	6434		6434	132	10	4	5	7	64	33			9
Gil-Bernal	2021	Spain	158	92	250		42	2					2				
Lebrun	2011	France	15220	5773	20993		820	4					4				
O'Connor	2014	Canada	755	339	1094		1094	30	5				25				
Polman	2006	Multi-country	660	282	942		627	5					3	1			1
Stamatellos	2021	Greece	129029	35529	164558		56767	1257					1257				

BCC: Basal cell carcinoma; LTC: leukemia (various types) and thyroid cancers; OCPC: Oral cavity and pharynx cancers; SCC: Squamous cell carcinoma; UN/NM: Unspecified Neoplasm/Not mentioned; FBC: Female breast cancer; BC: Breast cancer (male and female); CC: Cervical cancer; MM: Metastatic melanoma; CIN: Cervical Intraepithelial Neoplasia

Table 2: Quality assessment of included studies

Table X. Quality assessment checklist for included studies												
Author	Was the study's target population a close representation of the national population in relation to relevant variables, e.g., age, sex, occupation?	Was the sampling frame a true or close representation of the target population?	Was some form of random selection used to select the sample, OR, was a census undertaken?	Was the likelihood of non-response bias minimal?	Were data collected directly from subjects (as used in the study)?	Was an acceptable case definition used in the study? (e.g., prevalence of low back pain) shown to have reliability and validity (if necessary)?	Were the numerator (s) and denominator (s) for the parameter of interest appropriate	Was the same mode of data collection used for all subjects?	Were the numerator (s) and denominator (s) for the parameter of interest appropriate	Summary on the overall risk of study bias	Total score	
Alping	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk	1	
DAmico	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	0	
Foley	Low risk	Low risk	Low risk	High risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk	2	
Gil-Bernal	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	0	
Lebrun	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	0	
O'Connor	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk	1	
Polman	Low risk	Low risk	Low risk	Low risk	Low risk	High risk	Low risk	Low risk	Low risk	Low risk	1	
Stamatellos	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	Low risk	0	

References

1. Ghajarzadeh M, Jalilian R, Sahraian MA, Moghadasi AN, Azimi A, Mohammadifar M, *et al.* Pain in patients with multiple sclerosis. *Maedica* 2018;13:125.
2. Ghajarzadeh M, Azizi S, Moghadasi AN, Sahraian MA, Azimi A, Mohammadifar M, *et al.* Validity and reliability of the Persian version of the PERception de la Sclérose En Plaques et de ses Pousse'es questionnaire evaluating multiple sclerosis-related quality of life. *Int J Prev Med* 2016;7:25.
3. Piehl F. A changing treatment landscape for multiple sclerosis: Challenges and opportunities. *J Intern Med* 2014;275:364-81.
4. Alping P, Askling J, Burman J, Fink K, Fogdell-Hahn A, Gunnarsson M, *et al.* Cancer risk for fingolimod, natalizumab, and rituximab in multiple sclerosis patients. *Ann Neurol* 2020;87:688-99.
5. Rudick RA, Stuart WH, Calabresi PA, Confavreux C, Galetta SL, Radue EW, *et al.* Natalizumab plus interferon beta-1a for relapsing multiple sclerosis. *N Engl J Med* 2006;354:911-23.
6. Azimi A, Hanaei S, Sahraian MA, Mohammadifar M, Ramagopalan SV, Ghajarzadeh M. Incidence of seroconversion and sero-reversion in patients with multiple sclerosis (MS) who had been treated with natalizumab: A systematic review and meta-analysis. *J Clin Neurosci* 2020;71:129-34.
7. Hanaei S, Sahraian MA, Mohammadifar M, Ramagopalan SV, Ghajarzadeh M, Ghajarzadeh M. Prevalence of Anti-JC virus antibody seropositivity in patients with multiple sclerosis: A systematic review and meta-analysis. *Intervirology* 2019;62:169-73.
8. Brønnum-Hansen H, Koch-Henriksen N, Stenager E. Trends in survival and cause of death in Danish patients with multiple sclerosis. *Brain* 2004;127:844-50.
9. Grytten Torkildsen N, Lie S, Aarseth J, Nyland H, Myhr K. Survival and cause of death in multiple sclerosis: Results from a 50-year follow-up in Western Norway. *Mult Scler J* 2008;14:1191-8.
10. Hirst C, Swingler R, Compston D, Ben-Shlomo Y, Robertson NP. Survival and cause of death in multiple sclerosis: A prospective population-based study. *J Neurol Neurosurg Psychiatry* 2008;79:1016-21.
11. Smestad C, Sandvik L, Celius E. Excess mortality and cause of death in a cohort of Norwegian multiple sclerosis patients. *Mult Scler J* 2009;15:1263-70.
12. Ghajarzadeh M, Mohammadi A, Sahraian MA. Risk of cancer in multiple sclerosis (MS): A systematic review and meta-analysis. *Autoimmun Rev* 2020;19:102650.
13. Hoy D, Brooks P, Woolf A, Blyth F, March L, Bain C, *et al.* Assessing risk of bias in prevalence studies: Modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol* 2012;65:934-9.
14. Carbone ML, Lacal PM, Messinese S, De Giglio L, Pozzilli C, Persechino S, *et al.* Multiple sclerosis treatment and melanoma development. *Int J Mol Sci* 2020;21:2950.
15. Singer BA. The role of natalizumab in the treatment of multiple sclerosis: Benefits and risks. *Ther Adv Neurol Disord* 2017;10:327-36.
16. Lebrun C, Rocher F. Cancer risk in patients with multiple sclerosis: Potential impact of disease-modifying drugs. *CNS Drugs* 2018;32:939-49.
17. Bray F, Ren JS, Masuyer E, Ferlay J. Global estimates of cancer prevalence for 27 sites in the adult population in 2008. *Int J Cancer* 2013;132:1133-45.
18. Foley J, Carrillo-Infante C, Smith J, Evans K, Ho P-R, Lee L, *et al.* The 5-year Tysabri global observational program in safety (TYGRIS) study confirms the long-term safety profile of natalizumab treatment in multiple sclerosis. *Mult Scler Relat Disord* 2020;39:101863.
19. Stamatellos VP, Sifakis S, Papazisis G. Disease-modifying agents for multiple sclerosis and the risk for reporting cancer: A disproportionality analysis using US Food and Drug Administration Adverse Event Reporting System (FAERS) database. *Br J Clin Pharmacol* 2021;87:4769-79.
20. Polman CH, O'Connor PW, Havrdova E, Hutchinson M, Kappos L, Miller DH, *et al.* A randomized, placebo-controlled trial of natalizumab for relapsing multiple sclerosis. *N Engl J Med* 2006;354:899-910.
21. Babaesfahani A, Khanna NR, Kuns B. Natalizumab. *StatPearls*. Treasure Island (FL): StatPearls Publishing; 2021.
22. Castela E, Lebrun-Frenay C, Laffon M, Rocher F, Cohen M, Leccia NC, *et al.* Evolution of nevi during treatment with natalizumab: A prospective follow-up of patients treated with natalizumab for multiple sclerosis. *Arch Dermatol* 2011;147:72-6.