

Evaluation of Rubella IgG Antibodies Among Women at Marriage in Kermanshah City, Before and After Mass Vaccination

Hossein Hatami, Feizollah Mansouri¹, Yadollah Mehrabi², Hengameh Namdari³, Maryam Hatami⁴

Department of Public Health, ²Department of Epidemiology, School of Health, ⁴Department of Gynecology, Tajrish Shohada Hospital, Shahid Beheshti University of Medical Sciences, Tehran, ¹Department of Infectious Diseases, Kermanshah University of Medical Sciences Kermanshah, ³Epidemiology Executive of Center for Diseases Management, Tehran, Iran

Correspondence to:

Prof. Hossein Hatami,
Department of Public Health, School of Health, Shahid Beheshti University of Medical Sciences, Daneshjoo Blvd., Tabnak St., Evin, Tehran, Iran.
E-mail: hatami@hbi.ir

Date of Submission: Nov 11, 2011

Date of Acceptance: Apr 01, 2012

How to cite this article: Hatami H, Mansouri F, Mehrabi Y, Namdari H, Hatami M. Evaluation of rubella igg antibodies among women at marriage in kermanshah city, before and after mass vaccination. *Int J Prev Med* 2013;4:917-21.

ABSTRACT

Background: Rubella is a mild viral infectious disease, usually occurs subclinically without a serious complication. But if occurs during pregnancy, specially the first trimester, it can transmit and invade to fetus causing congenital rubella syndrome or fetal death. In 2003, nearly all Iranians at the age of 5 to 25 years old were vaccinated against rubella and the vaccine was included in the compulsory immunization, MMR afterwards.

Methods: This study was carried out on girls attended at a marriage consultancy center in Kermanshah city (West of Iran) in 2006. Considering 95% of confidence, the sample size was estimated at 140. All samples were tested using ELISA serologic method and compared with similar study conducted on 240 unvaccinated girls at the same place in 1999.

Results: Antibody positivity was observed in 99.3%, compared to 79.58% of the previous study. A rise of 19.72% (95% CI 12.7%-26.7%) of seropositivity was observed in vaccinated girls versus unvaccinated ones ($P < 0.001$).

Conclusion: In our previous study, we had recommended to screen for susceptibility to rubella before marriage, which is no longer required since more than 99% of vaccinated girls showed immunity at the time of marriage. However, as sustainability of immunity after rubella vaccination is usually less than immunity due to illness, we recommend screening for rubella protective antibody every few years.

Keywords: Congenital rubella, immunity, rubella, seroepidemiology, vaccination

INTRODUCTION

According to the WHO experts, if rubella infection occurs during the early stages of pregnancy, the potential of passing the virus through the placenta and reaching the fetus is about 90%.^[1] The most important complication of rubella is “congenital rubella syndrome (CRS),” which directly relates to lack of immunity during pregnancy. On the other hand, by vaccination of all the susceptible people or women in reproductive age, or the susceptible girls at the time of marriage and induce immunity

the immunity rate after rubella vaccination to be increased from 80% in 1999 to more than 90% in 2006, the sample size of 140 was calculated using the EpiInfo software and 95% confidence. All blood samples were taken and tested under similar conditions and by the same person at the reference laboratory. For quality assurance, about 10% of the samples were chosen, randomly, and rechecked at Red Crescent laboratory of Kermanshah. The titration was done by quantified ELISA method.

The data were analyzed and interpreted using percentages and confidence intervals (CI). The results of this study were compared to those of the previous similar study conducted by the main researcher in 1999 with similar design at the same place but unvaccinated girls.^[11] Comparison of proportions formula was used for this mean.

RESULTS

Age distribution indicated that the majority of the study population (35%) was 20–24 years old. Their average age was 21.95 (SD = 3.64), and the youngest and oldest were 15 and 30 years old, respectively. With respect to educational level, 20.14% of the samples were illiterate, 19.29% were in primary school, 20% in guidance school, 32.14% in secondary school, 5% have college degree, 11.43% were bachelor, and 2.86% were not identified. A sum of 19.29% of cases was living in rural and the rest (80.71%) in urban areas.

Positive antibody, immunity against rubella, was seen in 99.3% (139 out of 140 vaccinated girls, compared to 79.6% (191 out of 240 unvaccinated girls) [Figure 1] of the previous study. Therefore, we are faced to a significant rise of 19.7% (95% CI 12.7%–26.7%) of seropositivity and immunity against rubella in vaccinated versus unvaccinated girls ($Z = 5.481, P < 0.001$).

DISCUSSION

This study revealed that mass vaccination of rubella caused a highly significant decrease in the rate of negative antibody from 20.42% to 0.7%. The finding is in line with reliable sources^[7] and extensive country study, which has compared immunity rate before and after Iranian 2003 mass vaccination.^[14] The result also agrees with the research conducted in Mashhad (East of Iran).^[15] In addition, our study showed higher

immunity improvement than those reported from Shahre-Kord^[16] and Hamadan^[17] cities in the West of Iran.

During an extensive study made by Immunology Department researchers on 1217 serum samples before vaccination (on 324 susceptible and 893 immune persons), and 2007 serum samples after vaccination (24 susceptible and 1983 immune) in 5 to 15 years age group fulfilled in ten provinces, it was identified that the immunity rate of the prevaccination was raised from 73.4% to 98.8% of the postvaccination.^[14] In another study in Mashhad, the level of immunity before and after vaccination has been 70.38% and 98.5%, respectively.^[15] During a study made on 150 Shahr-e-Kord Medical University students, it was identified that 129 (86%) of them were immune before vaccination, while it increased to 96% after vaccination.^[16] During a study at Hamadan Health Center, the level of anti-rubella antibody in Hamadani girls at marriage had been 83.5%, while two years after such program it has increased to 98.4%^[17] [Table 1]. The effectiveness




Figure 1: Situation of anti-rubella antibody and immunity in unvaccinated (left) and vaccinated (right) Kermanshahi girls at marriage, 2006

Table 1: Rubella seropositivity before and after vaccination in Iran

Location	Vaccination		P Value
	Before	After	
Country ^[14]	73.40%	98.00%	$P < 0.001$
Mashhad ^[15]	70.38%	89.50%	$P < 0.001$
Shahr-e-Kord ^[16]	86.00%	96.90%	$P < 0.001$
Hamadan ^[17]	83.50%	98.40%	$P < 0.001$
Current study	79.58%	99.30%	$P < 0.001$

of the RA 27/3 vaccine has been demonstrated by the elimination of rubella and CRS from the western hemisphere and by the several European countries that have achieved and maintained high vaccination coverage with vaccines containing RA 27/3.^[18]

CONCLUSION

Mass rubella vaccination in 5 to 25 years old Iranian population during 2003,^[13] and continued thereafter as MMR in national vaccination program,^[19] has significantly resulted in increase in anti-rubella antibodies and immunity in high-risk people, as premarriage Kermanshahi girls, and in comparison to many of the results specified in local and international sources,^[7,20] and due to the susceptibility of less than 1% of the Kermanshahi girls at marriage, there is no more need to screen the vaccinated girls at marriage, which was emphasized in 1999;^[8] however, the screening studies is recommended every few years.

RECOMMENDATION

As the dispersed studies associated with the rubella seroepidemiology in different locations of the country resulted in awareness of lack of immunity of about 20% of the most vulnerable society members, i.e., the pregnant women, and after that the policy makers decided to announce the rubella vaccine as the compulsory state vaccines, it is proposed such studies to be made for the immunity status against tetanus, diphtheria, etc. Additionally, due to the fact that neither rubella infection nor vaccination result in life-long immunity, it is proposed that once each couple of years the studies associated with the rubella seroepidemiology to be performed in different regions of the country and if necessary, the vaccination of vulnerable society members be made, like in 2003.

ACKNOWLEDGEMENTS

The authors would like to thank the individuals who have been of tremendous assistance at the Rafatieh Health Centre and Kermanshah Province Health Department who cooperated in different steps of this research.

This study was financially supported by research deputy, Shahid Beheshti, University of Medical Sciences.

REFERENCES

1. Rubella and Congenital Rubella Syndrome (CRS), WHO, 2009. Available from: http://www.who.int/immunization_monitoring/diseases/rubella/en/index.html. [Last Cited on 2012 June 12].
2. Gershon AA. Rubella Virus (German Measles). In: Mandell, Douglass, Bennett's principles and practice of infectious diseases. 7th ed. USA: Elsevier Churchill Livingstone; 2010. p. 2127-32.
3. Park K. Rubella. In: Park's textbook of Preventive and Social Medicine. Delhi, India: Bhandot Publishers; 2009. p. 138-40.
4. Spiegel HM, Sever JL. Rubella virus. In: Gorbach SL, Bartlett JG, Blacklow NR. Infectious diseases. 3rd ed. Philadelphia: Lippincott; 2004. p. 2105-8.
5. Armstrong D, Cohen J. Infectious diseases. London: Mosby; 1999; 1 2.1.3-4/ 8.2.3-5
6. Cherry JD. Rubella virus. In: Feigin and Cherry's Text Book of Pediatric Infectious Diseases. 6th ed. Saunders; 2009. p. 2271-300.
7. Zimmerman LS, Reef SE. Rubella (German Measles). In: Fauci, Braunwald, Kasper editors. Harrison's Principles of Internal Medicine. 18th ed. New York: McGraw Hill; 2012. p. 1605-8.
8. Heymann DL. Control of communicable diseases manual. 19th ed. Washington, DC: American Public Health Association; 2008. p. 529-34.
9. Sullivan E, Burgess MA, Forrest GM. The epidemiology of Rubella and congenital rubella in Australia, 1992 to 1997. *Commun Dis Intell (CDI)* 1999;23:209-14.
10. De Haas R, Van den Hof S, Berbers GA, de Melker HE, Conyn-van Spaendonck MA. Prevalence of Antibodies against rubella Virus in the Netherlands 9 years after changing from selective to mass vaccination. *Epidemiol Infect* 1999;123:263-70.
11. Hatami H. Seroepidemiological Study of rubella in Girls at Marriage: Kermanshah. Iran, Iran, *J Ardabil Univ Med Sci* 2009;4:283-91.
12. Hatami H, Firoozi F. Sero-epidemiology of Rubella in Kermanshahi Girls at Marriage. *J Infect Dis Trop Med* 2002;7:65-71.
13. Esteghamati A, Gouya MM, Zahraei SM, Dadras MN, Rashidi A, Mahoney F. Progress in measles and Rubella elimination in Iran. *Pediatr Infect Dis J* 2007;26:1137-41.
14. Salimi H, Mokhtari Azad T, Tabatabaei H, Sarijloo M, Baethy K, Mahmoodi M, *et al.* Seroepidemiology of Rubella in 5 to 25 years old Iranian people, before and after 2003-mass vaccination in Iran. *J Infect Dis Trop Med* 2007;12:39-42.
15. Mahmoodi M, Vahedi E. A Comparative study of immunity against Rubella before and after 2003-mass vaccination. *J Mashhad Univ Med Sci* 2007;96:177-86.

16. Shirzad H, Shahabi G, Zamanzad B, Rafighian M. Efficacy of Rubella vaccine in students of Shahr-e-Kord University of Medical Sciences After Mass Vaccination Programm in Iran. *J Arak Univ Med Sci* 2005;8:30-7.
17. Eghbalian F, Ranjbar M. Seroepidemiology of Rubella in Hamadani girls at marriage, before and after mass vaccination programm. *J Iran Univ Med Sci* 2009;12:61-2.
18. WHO, Rubella vaccines: WHO position paper, *Weekly Epidemiological Record*, 2011;8629:301-16.
19. Expanded Program on Immunization (EPI), Islamic Republic of Iran Immunization Schedule, 7th ed. 2009.
20. WHO, Meeting of the Strategic Advisory Group of Experts on immunization, April 2012 – conclusions and recommendations, *Weekly Epidemiological Record*, 2012; 8721:201-16.

Source of Support: Nil **Conflict of Interest:** None declared.